

# Delta Waves for a Strongly Singular Initial-Boundary Hyperbolic Problem with Integral Boundary Condition

I. Kmit\*

22 July 2003

## Abstract

We investigate the existence and the singular structure of delta wave solutions to a semilinear strictly hyperbolic equation with strongly singular initial and boundary conditions. The boundary conditions are given in nonlocal form with a linear integral operator involved. We construct a delta wave solution as a distributional limit of solutions to the regularized system. This determines the macroscopic behavior of the corresponding generalized solution in the Colombeau algebra  $\mathcal{G}$  of generalized functions. We represent our delta wave as a sum of a purely singular part satisfying a linear system and a regular part satisfying a nonlinear system.

## 1 Introduction

In the domain

$$\Pi = \{(x, t) \in \mathbb{R}^2 \mid 0 < x < L, t > 0\}$$

we study the following initial-boundary value problem for the first-order semilinear hyperbolic equation:

$$(\partial_t + \lambda(x, t)\partial_x)u = p(x, t)u + f(x, t, u), \quad (x, t) \in \Pi \quad (1)$$

$$u|_{t=0} = a(x), \quad x \in (0, L) \quad (2)$$

$$u|_{x=0} = \int_0^L d(x, t)u \, dx, \quad t \in (0, \infty). \quad (3)$$

Mathematical models of this kind stem from mathematical biology and serve to describe the age-dependent population dynamics (see [2, 3, 12, 21, 22]). In particular,

---

\*Department of Numerical Mathematics & Programming, National University “Lvivska Polytechnika”, Bandera St. 12, 290646 Lviv, Ukraine. E-mail: [kmit@ov.litech.net](mailto:kmit@ov.litech.net) This work was done while visiting the Institut für Mathematik, Universität Wien, supported by an ÖAD grant.

the linear case of the problem, when  $f(x, t, u)$  does not depend on  $u$ , arises in demography, where  $u(x, t)$  is the population density of age  $x$  at time  $t$ ,  $a(x)$  is the initial density,  $d(x, t)$  is the birth rate,  $-p(x, t)$  is the death rate, and  $f(x, t)$  is the migrant density. Nonlinear models of age structured populations are studied in [2, 3]. To model point-concentration of the initial density and the birth rate, we consider the data  $a(x)$  and  $d(x, t)$  to be strongly singular, of the Dirac delta type.

As well known, solutions to the classical initial-boundary semilinear hyperbolic problems in a single space variable are at least as singular as the initial and the boundary data. We therefore can expect for the nonclassical problem (1)–(3) that the multiplication of distributions appears in the right-hand sides of (1) and (3). Such multiplication in general cannot be performed within the distributional theory and, by this reason, is usually defined in differential algebras of generalized functions. In [13] we used the Colombeau algebra of generalized functions  $\mathcal{G}(\overline{\Pi})$  [1, 4, 14] to prove a global existence-uniqueness result for (1)–(3). Nevertheless, the macroscopic behavior of the Colombeau solutions remained unclear.

We here show that the Colombeau solution to (1)–(3) is associated to the distributional solution. This means that the system has a delta wave solution in the sense of [20], i.e., the sequence of approximate (or sequential) solutions obtained by regularizing all singular data has a weak limit. In the course of construction of the delta wave solution we show interaction and propagation of singularities.

It should be noted that an associated distribution or in other terminology, a delta wave solution, though contains an important information about the singular structure of the generalized solution, in general does not satisfy the system in a differential-algebraic sense. Our paper brings one more example into the collection of associated distributions which are not distributional solutions.

The advantage of using delta wave solutions lies in the fact that, due to the procedure of their obtaining, they are always stable. In contrast with this, if we use a priori defined intrinsic multiplication of distributions for obtaining distributional solutions, the result may be nonstable and noncorrect [5]. The concept of a delta wave solution has also other advantages. It allows us to solve nonlinear systems and systems with nonsmooth coefficients, for which the distributional theory is not well adapted. For the delta wave solutions of semilinear hyperbolic problems we refer the reader to the sources [5, 6, 9] and [13]–[20].

We split a delta wave into the sum of a regular part satisfying a nonlinear equation and a singular part satisfying a linear equation. The idea of nonlinear splitting goes back to [16, 20, 17, 6]. An important feature of the nonlinear splitting suggested here is quite strong interdependence of the singular and the regular parts. A similar phenomenon is discovered in [13] for a nonlocal problem with nonseparable boundary conditions, where the singular part of the nonlinear splitting depends on the regular part.

Delta wave solutions for initial-boundary semilinear hyperbolic problems were considered in [18, 13]. The paper [18] investigates the existence and structure of delta waves in a nonlinear boundary value problem for a second order hyperbolic equation where the boundary condition is nonlinear and the nonlinearity is given by a bounded smooth function. Both in [18] and [13] the right hand side of the

differential equations is bounded and in [18] it can be also sublinear with respect to  $u$ .

The paper is organized as follows. In Section 2 we describe in detail our splitting of a delta wave solution and state our main result. The proof is given in Sections 3–8. In particular, in Section 5 we show that our splitting procedure is correct. In Section 6 we are concerned with the regular part. Using the Cauchy criterion of the uniform convergency, we prove that the family of approximate solutions to the regular part uniformly converges on any compact subset of  $\overline{\Pi}$ . In Section 7 we deal with the singular part and prove that the sequence of approximate solutions to the singular part converges in  $\mathcal{D}'(\overline{\Pi})$  to a function  $v$ . We then show that  $v$  actually represents the purely singular part of the initial problem and is the sum of measures concentrated on characteristic curves (see Sections 7 and 8).

## 2 Interaction and propagation of strong singularities and construction of a delta wave solution

We first list assumptions that will be made for the problem (1)–(3).

*Assumption 1.*  $a(x) = a_s(x) + a_r(x)$ ,  $d(x, t) = b(x) \otimes c(t) = (b_s(x) + b_r(x)) \otimes (c_s(t) + c_r(t))$ , where  $a_s(x)$ ,  $b_s(x)$ ,  $c_s(t)$  and  $a_r(x)$ ,  $b_r(x)$ ,  $c_r(t)$  are, respectively, singular and regular parts of the functions  $a(x)$ ,  $b(x)$ , and  $c(t)$ .

*Assumption 2.*  $a_s(x)$ ,  $b_s(x)$ , and  $c_s(t)$  are the finite sums of the delta functions at points, whose supports are as follows:

$\text{supp } a_s(x) = \{x_1^*, x_2^*, \dots, x_m^*\}$ , where  $0 < x_1^* < \dots < x_m^* < L$ .

$\text{supp } b_s(x) = \{x_1, x_2, \dots, x_k\}$ , where  $0 < x_1 < \dots < x_k < L$ .

$\text{supp } c_s(t) = \{t_1, t_2, \dots, t_l\}$ , where  $0 < t_1 < \dots < t_l$ .

*Assumption 3.*  $b_r(0) = 0$  and  $b_r(L) = 0$ .

*Assumption 4.*  $a_r(0) = 0$  and  $c_r(0) = 0$ .

*Assumption 5.*  $p, f, a_r, b_r, c_r$  are continuous and  $\lambda$  is continuously differentiable with respect to all their arguments,  $f$  is continuously differentiable with respect to  $u$ .

*Assumption 6.*  $\lambda(x, t) > 0$  for all  $(x, t) \in \overline{\Pi}$ .


*Assumption 7.*  $f$  and  $\nabla_u f$  are globally bounded with respect to  $(x, t)$  varying in compact subsets of  $\overline{\Pi}$ .

Assumption 4 serves to ensure the 0-order compatibility between (2) and (3). Note that the assumptions are not restrictive from the viewpoint of applications.

Recall that all characteristics of the differential equation (1) are solutions to the following initial problem for ordinary differential equation:

$$\frac{d\xi}{d\tau} = \lambda(\xi(\tau), \tau), \quad \xi(t) = x,$$

where  $(x, t) \in \overline{\Pi}$ . It is well known that, under Assumptions 5 and 6, for every  $(x, t) \in \overline{\Pi}$  this problem has a unique  $C^1$ -solution which can be expressed in any of



two forms  $\xi = \omega(\tau; x, t)$  or  $\tau = \tilde{\omega}(\xi; x, t)$ .

Choose  $\varepsilon_0 > 0$  so small that  $x_1^* - \varepsilon_0 > 0$  and  $t_1 - \varepsilon_0 > 0$ . Some additional conditions on  $\varepsilon_0$  will be put below. We will consider  $\varepsilon$  in the range  $0 < \varepsilon \leq \varepsilon_0$ .

**Definition 1** Let  $I_-$  be the union of the characteristics  $\omega(t; x_i, t_j)$  for all  $i \leq k$  and  $j \leq l$ . Let  $I_-^\varepsilon$  be the union of the neighborhoods  $\{(\omega(\tau; x_i, t), \tau) \mid \tilde{\omega}(x_i; x_i - \varepsilon, t_j + \varepsilon) < t < \tilde{\omega}(x_i; x_i + \varepsilon, t_j - \varepsilon)\}$  for all  $i \leq k$  and  $j \leq l$ .

**Definition 2** Let  $I_+ = \bigcup_{n \geq 0} I_+[n]$  and, for  $\varepsilon < \varepsilon_0$ ,  $I_+^\varepsilon = \bigcup_{n \geq 0} I_+^\varepsilon[n]$ , where  $I_+[n]$  and  $I_+^\varepsilon[n]$  are subsets of  $\overline{\Pi}$  defined by induction as follows.

- $I_+[0]$  includes the characteristics  $\omega(t; x_i^*, 0)$  and  $\omega(t; 0, t_j)$  for all  $i \leq m$  and  $j \leq l$  (i.e.  $I_+[0]$  is the union of these characteristics).

$I_+^\varepsilon[0]$  includes the neighborhoods  $\{(\omega(\tau; x, 0), \tau) \mid x_i^* - \varepsilon < x < x_i^* + \varepsilon\}$  and  $\{(\omega(\tau; 0, t), \tau) \mid t_j - \varepsilon < t < t_j + \varepsilon\}$ .

- Let  $n \geq 1$ . If  $I_+[n-1]$  includes the characteristic  $\omega(t; x_i, \tilde{t})$ , then  $I_+[n]$  includes the characteristic  $\omega(t; 0, \tilde{t})$ .

If  $I_+^\varepsilon[n-1]$  includes the neighborhood  $\{(\omega(\tau; x_i, t), \tau) \mid \tilde{t} - \varepsilon^- < t < \tilde{t} + \varepsilon^+\}$ , then  $I_+^\varepsilon[n]$  includes the neighborhood  $\{(\omega(\tau; 0, t), \tau) \mid \tilde{\omega}(x_i - \varepsilon; x_i, \tilde{t} - \varepsilon^-) < t < \tilde{\omega}(x_i + \varepsilon; x_i, \tilde{t} + \varepsilon^+)\}$ .

The set  $I_+$  captures the propagation of all singularities. For characteristics contributing into  $I_+$  (respectively,  $I_+ \setminus I_+[0]$ ), denote their intersection points with the axis  $x = 0$  by  $t_1^*, t_2^*, \dots$  (respectively,  $\tilde{t}_{i_1}, \tilde{t}_{i_2}, \dots$ ). We assume that  $t_j^* < t_{j+1}^*$  for  $j \geq 1$  and  $i_n = p$  for  $\tilde{t}_{i_n} = t_p^*$ . Obviously,  $\{t_1^*, t_2^*, \dots\} = \{t_1, \dots, t_l\} \cup \{\tilde{t}_{i_1}, \tilde{t}_{i_2}, \dots\}$ . Let  $\varepsilon_i^-(\varepsilon)$  and  $\varepsilon_i^+(\varepsilon)$  be such that  $I_+^\varepsilon \cap \{(x, t) \in \bar{\Pi} \mid x = 0\} = \bigcup_i \{(0, t) \mid t_i^* - \varepsilon_i^-(\varepsilon) < t < t_i^* + \varepsilon_i^+(\varepsilon)\}$ . If  $t_i^* = t_j$  for some  $j \leq l$ , then  $\varepsilon_i^-(\varepsilon) = \varepsilon_i^+(\varepsilon) = \varepsilon$ . Observe that  $\lim_{\varepsilon \rightarrow 0} \varepsilon_i^-(\varepsilon) = 0$  and  $\lim_{\varepsilon \rightarrow 0} \varepsilon_i^+(\varepsilon) = 0$ .

*Assumption 8.*  $\tilde{\omega}(0; x_i, t_j) \neq t_s^*$ ,  $\omega(0; x_i, t_j) \neq x_q^*$  for all  $i \leq k$ ,  $j \leq l$ ,  $q \leq m$  and  $t_s^* < t_l$ .

This assumption means that no three different singularities caused by the initial and the boundary data hit at the same point. In other words, neither points  $(x_q^*, 0)$  and  $(x_i, t_j)$  nor points  $(0, t_s^*)$  and  $(x_i, t_j)$  are connected by any of characteristic curves. As a consequence, there exists  $\varepsilon_0$  such that, for each  $\varepsilon \leq \varepsilon_0$ ,  $I_-^\varepsilon \cap I_+^\varepsilon = \emptyset$ . Assume that  $(0, 0) \notin I_-$ . We choose  $\varepsilon_0$  so small that  $I_-^{\varepsilon_0}$  and  $I_+^{\varepsilon_0}$  do not contain the point  $(0, 0)$ . Clearly,  $\bigcap_{\varepsilon > 0} I_+^\varepsilon = I_+$  and  $\bigcap_{\varepsilon > 0} I_-^\varepsilon = I_-$ .

Our aim is to show that the generalized solution to the problem (1)–(3), whose existence is shown in [13], admits an *associated distribution* or a *delta-wave*. The latter means that the family  $(u^\varepsilon)_{\varepsilon > 0}$  of solutions to the system with regularized initial and boundary data

$$(\partial_t + \lambda(x, t)\partial_x)u^\varepsilon = p(x, t)u^\varepsilon + f(x, t, u^\varepsilon) \quad (4)$$

$$u^\varepsilon|_{t=0} = a_s^\varepsilon + a_r \quad (5)$$

$$u^\varepsilon|_{x=0} = (c_s^\varepsilon + c_r) \int_0^L (b_s^\varepsilon + b_r)u^\varepsilon dx \quad (6)$$

has a weak limit. Here

$$a_s^\varepsilon = a_s * \varphi_\varepsilon, \quad b_s^\varepsilon = b_s * \varphi_\varepsilon, \quad c_s^\varepsilon = c_s * \varphi_\varepsilon,$$

where mollifiers  $\varphi_\varepsilon$  are model delta nets, that is,

$$\varphi_\varepsilon(x) = \frac{1}{\varepsilon} \varphi\left(\frac{x}{\varepsilon}\right)$$

for an arbitrary fixed  $\varphi \in \mathcal{D}(\mathbb{R})$  with  $\int \varphi(x) dx = 1$ . Note that

$$a_s^\varepsilon = O\left(\frac{1}{\varepsilon}\right), \quad b_s^\varepsilon = O\left(\frac{1}{\varepsilon}\right), \quad c_s^\varepsilon = O\left(\frac{1}{\varepsilon}\right) \quad (7)$$

and

$$\int_0^L |a_s^\varepsilon(x)| dx \leq C, \quad \int_0^L |b_s^\varepsilon(x)| dx \leq C, \quad \int_0^\infty |c_s^\varepsilon(t)| dt \leq C, \quad (8)$$

where  $C$  does not depend on  $\varepsilon$ . We will consider mollifiers  $\varphi$  with

$$\text{supp } \varphi \subset [-1, 1]. \quad (9)$$

This restriction makes no loss of generality, because if (9) is not true, then  $\text{supp } \varphi \subset [-d, d]$  for some  $d > 0$ . Therefore  $\text{supp } \varphi_\varepsilon \subset [-d\varepsilon, d\varepsilon]$  and it is enough to replace  $I_+^\varepsilon$  by  $I_+^{d\varepsilon}$  to keep all arguments valid, with the result not depending on  $d$ .

It follows from (9) that for all  $\varepsilon > 0$

$$\begin{aligned} \int_{x_i^* - \varepsilon}^{x_i^* + \varepsilon} a_s^\varepsilon(x) dx &= 1, \quad 1 \leq i \leq m; & \int_{x_j - \varepsilon}^{x_j + \varepsilon} b_s^\varepsilon(x) dx &= 1, \quad 1 \leq j \leq k; \\ \int_{t_p - \varepsilon}^{t_p + \varepsilon} c_s^\varepsilon(t) dt &= 1, \quad 1 \leq p \leq l. \end{aligned} \quad (10)$$

Let  $T$  be an arbitrary positive real,  $\Pi^T = \{(x, t) \in \Pi \mid t < T\}$ . We will show that a delta-wave splits up into the sum  $w + v$  of the following kind.

The function  $w$  corresponds to the regular part of the problem. More specifically, for every  $T > 0$ , the restriction of  $w$  to  $\Pi^T$  is the limit of  $w^\varepsilon$  in  $C(\Pi^T)$  as  $\varepsilon \rightarrow 0$ , where  $w^\varepsilon$  for every fixed  $\varepsilon > 0$  is a continuous solution to the nonlinear problem

$$(\partial_t + \lambda(x, t)\partial_x)w^\varepsilon = p(x, t)w^\varepsilon + f(x, t, w^\varepsilon) \quad (11)$$

$$w^\varepsilon|_{t=0} = a_r \quad (12)$$

$$w^\varepsilon|_{x=0} = c_r \int_0^L [(b_s^\varepsilon + b_r)w^\varepsilon + b_r v^\varepsilon] dx. \quad (13)$$

The function  $v$  corresponds to the singular part of the problem and is the limit of  $v^\varepsilon$  in  $\mathcal{D}'(\Pi)$  as  $\varepsilon \rightarrow 0$ , where  $v^\varepsilon$  for every fixed  $\varepsilon > 0$  is a continuous solution to the linear problem

$$(\partial_t + \lambda(x, t)\partial_x)v^\varepsilon = p(x, t)v^\varepsilon \quad (14)$$

$$v^\varepsilon|_{t=0} = a_s^\varepsilon \quad (15)$$

$$v^\varepsilon|_{x=0} = c_s^\varepsilon \int_0^L [(b_s^\varepsilon + b_r)w^\varepsilon + b_r v^\varepsilon] dx + c_r \int_0^L b_s^\varepsilon v^\varepsilon dx. \quad (16)$$

**Proposition 1** *For every  $\varepsilon \leq \varepsilon_0$  there exist a unique  $C(\overline{\Pi})$ -solution  $u^\varepsilon$  to the problem (4)–(6), a unique  $C(\overline{\Pi})$ -solution  $w^\varepsilon$  to the problem (11)–(13), and a unique  $C(\overline{\Pi})$ -solution  $v^\varepsilon$  to the problem (14)–(16).*

*Proof.* For every fixed  $\varepsilon > 0$ , (1)–(3) is a special case of the problem studied in [13]. From the proof of [13, Theorem 3] it follows that, if Assumptions 4–7 hold and  $\varepsilon$  is so small that  $(0, 0) \notin I_+^\varepsilon$ , this problem has a unique  $C(\overline{\Pi})$ -solution  $u^\varepsilon$ .

Fix  $\varepsilon > 0$ . We consider (11)–(13) and (14)–(16) simultaneously thereby obtaining an initial-boundary value problem for a system of two hyperbolic equations with respect to  $(w^\varepsilon, v^\varepsilon)$ . This is another special case of the problem studied in [13]. Note that we have zero-order compatibility of (12), (13) and of (15), (16), the former by Assumption 4 and the latter by Assumption 4 and the fact that  $(0, 0) \notin I_+^\varepsilon$ . From the proof of [13, Theorem 3] it follows that under Assumptions 4–7 the problem has a unique  $C(\overline{\Pi})$ -solution  $(w^\varepsilon, v^\varepsilon)$ .  $\square$

We are now prepared to state the main result of the paper.

**Theorem 1** *Let Assumptions 1–8 hold. Let  $u^\varepsilon$ , for every  $\varepsilon > 0$ , be the continuous solution to the problem (4)–(6). Then*

$$u^\varepsilon \rightarrow w + v \quad \text{in } \mathcal{D}'(\Pi) \quad \text{as } \varepsilon \rightarrow 0,$$

where

- for every  $T > 0$ , the restriction of  $w$  to  $\Pi^T$  is the limit of  $w^\varepsilon$  in  $C(\Pi^T)$  as  $\varepsilon \rightarrow 0$  with  $w^\varepsilon$  being the continuous solution to the problem (11)–(13),
- $v = \lim_{\varepsilon \rightarrow 0} v^\varepsilon$  in  $\mathcal{D}'(\Pi)$  with  $v^\varepsilon$  being the continuous solution to the problem (14)–(16). Furthermore, the restriction of  $v$  to  $\Pi \setminus I_+$  is identically equal to 0.

**Corollary**  $\text{sing supp}(w + v) = \text{sing supp } v = \text{supp } v \subset I_+$ .

This means that  $v$  actually represents the purely singular part of the initial problem. The proof of the corollary is straightforward.

The proof of Theorem 1 consists of five lemmas whose proofs are given in Sections 4–8.

**Lemma 1** *Let Assumptions 1, 2, 4–6, and 8 hold and  $v^\varepsilon$  be as in Theorem 1. Then*

$$v^\varepsilon \rightarrow 0 \quad \text{pointwise off } I_+ \quad \text{as } \varepsilon \rightarrow 0.$$

**Lemma 2** *Let Assumptions 1–8 hold and  $u^\varepsilon, v^\varepsilon$ , and  $w^\varepsilon$  be as in Theorem 1. Then*

$$u^\varepsilon - v^\varepsilon - w^\varepsilon \rightarrow 0 \quad \text{in } L_{loc}^1(\Pi) \quad \text{as } \varepsilon \rightarrow 0.$$

**Lemma 3** *Let Assumptions 1–8 hold and  $w^\varepsilon$  be as in Theorem 1. Then*

$$w^\varepsilon \quad \text{converges in } C(\overline{\Pi^T}) \quad \text{as } \varepsilon \rightarrow 0$$

for an arbitrary fixed  $T > 0$ .

**Lemma 4** *Let Assumptions 1–8 hold and  $v^\varepsilon$  be as in Theorem 1. Then*

$$v^\varepsilon \quad \text{converges in } \mathcal{D}'(\Pi) \quad \text{as } \varepsilon \rightarrow 0.$$

**Lemma 5** *Let Assumptions 1–8 hold,  $v^\varepsilon$  be as in Theorem 1, and  $v = \lim_{\varepsilon \rightarrow 0} v^\varepsilon$  in  $\mathcal{D}'(\Pi)$ . Then  $v$  restricted to  $\Pi \setminus I_+$  is identically equal to 0.*

Theorem 1 now follows from the embedding of  $L_{loc}^1(\Pi)$  into  $\mathcal{D}'(\Pi)$ .

### 3 Representation of the problems (11)–(13) and (14)–(16) in an integral-operator form

The problem (11)–(13) is equivalent to the integral-operator equation

$$w^\varepsilon(x, t) = (Rw^\varepsilon)(x, t) + \int_{\theta(x, t)}^t \left[ f(\xi, \tau, w^\varepsilon) + (pw^\varepsilon)(\xi, \tau) \right] \Big|_{\xi=\omega(\tau; x, t)} d\tau \quad (17)$$

and to the corresponding linearized integral-operator equation

$$\begin{aligned} w^\varepsilon(x, t) &= (Rw^\varepsilon)(x, t) \\ &+ \int_{\theta(x, t)}^t \left[ f(\xi, \tau, 0) + w^\varepsilon \left( \int_0^1 (\nabla_u f)(\xi, \tau, \sigma w^\varepsilon) d\sigma + p(\xi, \tau) \right) \right] \Big|_{\xi=\omega(\tau; x, t)} d\tau \end{aligned} \quad (18)$$

with boundary operator

$$(Rw^\varepsilon)(x, t) = \begin{cases} a_r(\omega(0; x, t)) & \text{if } \theta(x, t) = 0, \\ w^\varepsilon(0, \theta(x, t)) & \text{if } \theta(x, t) > 0. \end{cases} \quad (19)$$

Here

$$\theta(x, t) = \min_{(\omega(\tau; x, t), \tau) \in \partial\Pi} \tau.$$

The boundary function  $w^\varepsilon(0, t)$  is given by (13).

The problem (14)–(16) is equivalent to the integral-operator equation

$$v^\varepsilon(x, t) = (Rv^\varepsilon)(x, t) + \int_{\theta(x, t)}^t (pv^\varepsilon)(\omega(\tau; x, t), \tau) d\tau \quad (20)$$

with boundary operator

$$(Rv^\varepsilon)(x, t) = \begin{cases} a_s^\varepsilon(\omega(0; x, t)) & \text{if } \theta(x, t) = 0, \\ v^\varepsilon(0, \theta(x, t)) & \text{if } \theta(x, t) > 0. \end{cases} \quad (21)$$

The function  $v^\varepsilon(0, t)$  is given by the formula (16). The continuous solution of (20) can be expressed in the form

$$v^\varepsilon(x, t) = S(x, t)(Rv^\varepsilon)(x, t) \quad (22)$$

with  $C(\overline{\Pi})$ -function

$$\begin{aligned} S(x, t) &= 1 + \int_{\theta(x, t)}^t p(\omega(\tau; x, t), \tau) d\tau \\ &+ \int_{\theta(x, t)}^t p(\omega(\tau; x, t), \tau) d\tau \int_{\theta(x, t)}^\tau p(\omega(\tau_1; \omega(\tau; x, t), \tau), \tau_1) d\tau_1 + \dots \end{aligned} \quad (23)$$



## 4 Proof of Lemma 1

By (22) it suffices to show for every  $(x, t) \in \Pi \setminus I_+$  that, if  $\varepsilon$  is small enough, then  $(Rv^\varepsilon)(x, t) = 0$ . If  $\theta(x, t) = 0$ , the latter is true by the equality  $(Rv^\varepsilon)(x, t) = a_s^\varepsilon(\omega(0; x, t))$  and the fact that  $(\omega(0; x, t), 0) \notin I_+$ . Consider the case that  $\theta(x, t) > 0$ . Since  $\theta(x, t) \notin (\overline{I_+^\varepsilon} \cap \{(x, t) \mid x = 0\})$ , the proof will be complete by showing that

$$\text{supp } v^\varepsilon(0, t) \subset (\overline{I_+^\varepsilon} \cap \{(x, t) \mid x = 0\}), \quad (24)$$

where  $v^\varepsilon(0, t)$  is defined by (16). Observe that (24) is true for the first summand in (16). Indeed, by (9), Assumption 2, and the definition of  $I_+^\varepsilon$ ,

$$\begin{aligned} \text{supp} \left( c_s^\varepsilon \int_0^L [(b_s^\varepsilon + b_r)w^\varepsilon + b_r v^\varepsilon] dx \right) &\subset \bigcup_{i=1}^l [t_i - \varepsilon, t_i + \varepsilon] \subset \text{supp } c_s^\varepsilon \\ &\subset (\overline{I_+^\varepsilon[0]} \cap \{(x, t) \mid x = 0\}). \end{aligned} \quad (25)$$

To obtain (24), for the second summand in (16) we prove the inclusion

$$\text{supp} \left( c_r \int_0^L b_s^\varepsilon v^\varepsilon dx \right) \subset \bigcup_{n \geq 1} [\tilde{t}_{i_n} - \varepsilon_{i_n}^-(\varepsilon), \tilde{t}_{i_n} + \varepsilon_{i_n}^+(\varepsilon)]. \quad (26)$$

Recall that  $\tilde{t}_{i_n}$ ,  $n \geq 1$ , are intersection points of  $I_+ \setminus I_+[0]$  with the axis  $x = 0$ . Suppose (26) is false. Then there exists

$$t^1 \notin \bigcup_{n \geq 1} [\tilde{t}_{i_n} - \varepsilon_{i_n}^-(\varepsilon), \tilde{t}_{i_n} + \varepsilon_{i_n}^+(\varepsilon)] \quad (27)$$

such that

$$\int_0^L b_s^\varepsilon(x) v^\varepsilon(x, t^1) dx = \int_0^L b_s^\varepsilon(x) (Rv^\varepsilon)(x, t^1) S(x, t^1) dx \neq 0. \quad (28)$$

We fix such  $t^1$  and set

$$J^1 = \text{supp } b_s^\varepsilon(x) \cap \text{supp } v^\varepsilon(x, t^1). \quad (29)$$

By (28)

$$\text{mes } J^1 \neq 0. \quad (30)$$

Assume that  $\theta(x_0, t^1) = 0$  for some  $x_0 \in J^1$ . By (21) and (29),

$$(Rv^\varepsilon)(x_0, t^1) = a_s^\varepsilon(\omega(0; x_0, t^1)) \neq 0.$$

This means that  $\omega(0; x_0, t^1) \in \text{supp } a_s^\varepsilon \subset \overline{I_+^\varepsilon} \cap \{(x, t) \mid x = 0\}$ . We conclude that  $(x_0, t^1) \in \overline{I_+^\varepsilon[0]}$ . Furthermore, from (29) we have  $x_0 \in [x_i - \varepsilon, x_i + \varepsilon]$  for some  $i \leq k$ . From the definition of  $I_+^\varepsilon$  it follows that, if  $(x, t) \in \overline{I_+^\varepsilon[j]}$  and  $x \in [x_i - \varepsilon, x_i + \varepsilon]$  for some  $i \leq k$ , then  $(0, t) \in \overline{I_+^\varepsilon[j+1]}$ . Hence  $(0, t^1) \in \overline{I_+^\varepsilon[1]}$ . This contradicts (27).

Assume therefore that  $\theta(x, t^1) > 0$  for all  $x \in J^1$ . Then in (28) we have  $(Rv^\varepsilon)(x, t^1) = v^\varepsilon(0, \theta(x, t^1))$  and therefore

$$\int_0^L b_s^\varepsilon(x) v^\varepsilon(0, \theta(x, t^1)) S(x, t^1) dx \neq 0.$$

By (29) and (30) there exists  $t^2 \in \theta(J^1, t^1)$  such that  $v^\varepsilon(0, t^2) \neq 0$ . It is clear that  $t^2 < t^1$ . Assume that  $(0, t^2) \in \overline{I_+^\varepsilon}$ . Let  $x_0$  be such that  $\theta(x_0, t^1) = t^2$ . By the definition of  $I_+^\varepsilon$ ,  $(x_0, t^1) \in \overline{I_+^\varepsilon[j]}$  for some  $j \geq 0$ . Furthermore,  $x_0 \in J^1$  and therefore  $x_0 \in [x_i - \varepsilon, x_i + \varepsilon]$  for some  $i \leq k$ . Hence  $(0, t^1) \in \overline{I_+^\varepsilon[j+1]}$ . This again contradicts (27).

Assume therefore that  $(0, t^2) \notin \overline{I_+^\varepsilon}$ . On the account of (16) and (25), we rewrite the condition  $v^\varepsilon(0, t^2) \neq 0$  as

$$\int_0^L b_s^\varepsilon(x) v^\varepsilon(x, t^2) dx \neq 0.$$

Set

$$J^2 = \text{supp } b_s^\varepsilon(x) \cap \text{supp } v^\varepsilon(x, t^2).$$

Note that  $\text{mes } J^2 \neq 0$ . Similarly to the above, if  $\theta(x_0, t^2) = 0$  for some  $x_0 \in J^2$ , then  $(0, t^2) \notin \overline{I_+^\varepsilon[1]}$ , a contradiction with (27). We therefore assume that  $\theta(\xi, t^2) > 0$  for all  $x \in J^2$  and continue in this fashion, thereby constructing sequences  $t^k \in \theta(I^{k-1}, t^{k-1})$  and  $J^k = \text{supp } b_s^\varepsilon(x) \cap \text{supp } v^\varepsilon(x, t^k)$  for  $k \geq 2$  such that  $v^\varepsilon(0, t^k) \neq 0$  and  $(0, t^k) \notin \overline{I_+^\varepsilon}$ . By Assumptions 5 and 6, for some

$$k \leq \left\lceil \frac{T \max_{(x,t) \in \overline{\Pi^T}} |\lambda|}{x_1 - \varepsilon_0} \right\rceil$$

there exists  $x_0 \in J^k$  such that  $\theta(x_0, t^k) = 0$ . This implies  $(0, t^k) \in \overline{I_+^\varepsilon[1]}$ , a contradiction with (27).

Thus (26) is true and the proof of Lemma 1 is complete.

For the further reference observe that

$$\text{supp } v^\varepsilon \subset \overline{I_+^\varepsilon}. \quad (31)$$

This fact is true by (21), (22), (24), and the definition of  $I_+^\varepsilon$ .

## 5 Proof of Lemma 2

Choose  $\varepsilon_0$  so small that the number of connected components of  $\Pi^T \cap I_+^{\varepsilon_0}$  and  $\Pi^T \cap I_+$  coincide.

**Definition 3** Given  $T > 0$ , let  $\Pi_0^T = \{(x, t) \in \Pi^T \mid \omega(t; 0, 0) < x\}$  and  $\Pi_1^T = \Pi^T \setminus \overline{\Pi_0^T}$ .

Let  $n(T)$  and  $\rho(T)$  be the number of connected components of  $\Pi_1^T \setminus \overline{I_+^\varepsilon}$  (and  $\Pi_1^T \setminus I_+$ ) and  $\Pi_1^T \cap I_+^\varepsilon$  (and  $\Pi_1^T \cap I_+$ ), respectively. We denote these components, respectively, by  $\Pi^\varepsilon(1), \dots, \Pi^\varepsilon(n(T))$  ( $\Pi(1), \dots, \Pi(n(T))$ ) and  $I_+^\varepsilon(1), \dots, I_+^\varepsilon(\rho(T))$  ( $I_+(1), \dots, I_+(\rho(T))$ ).

Clearly,  $\Pi(i) = \bigcup_{\varepsilon>0} \Pi^\varepsilon(i)$  and  $I_+(i) = \bigcap_{\varepsilon>0} I_+^\varepsilon(i)$ . Observe that  $\rho(T)$  does not depend on  $\varepsilon$  and either  $n(T) = \rho(T)$  or  $n(T) = \rho(T) + 1$ . In the latter case, if  $n(T) = \rho(T) + 1$ , we define  $I_+^\varepsilon(\rho(T) + 1) = \emptyset$ .

Given  $T$ , we choose  $\varepsilon_0$  so small that for all  $\varepsilon \leq \varepsilon_0$

$$\text{supp } b_s^\varepsilon \subset (\omega(t_i^* + \varepsilon_i^+(\varepsilon); 0, t_i^* - \varepsilon_i^-(\varepsilon)), L]. \quad (32)$$

From (4)–(6), (11)–(13), and (14)–(16) it follows that the difference  $u^\varepsilon - v^\varepsilon - w^\varepsilon$  satisfies the system

$$\begin{aligned} (\partial_t + \lambda(x, t)\partial_x)(u^\varepsilon - v^\varepsilon - w^\varepsilon) &= p(x, t)(u^\varepsilon - v^\varepsilon - w^\varepsilon) \\ &+ F(x, t)(u^\varepsilon - v^\varepsilon - w^\varepsilon) + f(x, t, u^\varepsilon) - f(x, t, u^\varepsilon - v^\varepsilon), \end{aligned} \quad (33)$$

$$(u^\varepsilon - v^\varepsilon - w^\varepsilon)|_{t=0} = 0, \quad (34)$$

$$\begin{aligned} (u^\varepsilon - v^\varepsilon - w^\varepsilon)|_{x=0} &= c_s^\varepsilon \int_0^L b_s^\varepsilon(u^\varepsilon - w^\varepsilon) dx \\ &+ (c_s^\varepsilon + c_r) \int_0^L b_r(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx + c_r \int_0^L b_s^\varepsilon(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx, \end{aligned} \quad (35)$$

where

$$F(x, t) = \int_0^1 (\nabla_u f)(x, t, \sigma(u^\varepsilon - v^\varepsilon) + (1 - \sigma)w^\varepsilon) d\sigma.$$

**Claim 1**  $u^\varepsilon - v^\varepsilon - w^\varepsilon \rightarrow 0$  in  $L^1(\overline{\Pi_0^T})$  as  $\varepsilon \rightarrow 0$ .

*Proof.* The problem (33)–(35) on  $\Pi_0^T$  reduces to the Cauchy problem (33)–(34). By Assumption 7,  $f$  is globally bounded and, by Lemma 1,  $f(x, t, u^\varepsilon) - f(x, t, u^\varepsilon - v^\varepsilon) \rightarrow 0$  as  $\varepsilon \rightarrow 0$  pointwise off  $I_+$ . By Lebesgue's dominated theorem,  $f(x, t, u^\varepsilon) - f(x, t, u^\varepsilon - v^\varepsilon) \rightarrow 0$  in  $L^1(\overline{\Pi_0^T})$  as  $\varepsilon \rightarrow 0$ . Applying Lebesgue's dominated theorem to the functions  $u^\varepsilon - v^\varepsilon - w^\varepsilon$  defined by (33)–(34), we obtain the claim.  $\square$

**Claim 2**  $u^\varepsilon - v^\varepsilon - w^\varepsilon \rightarrow 0$  pointwise for  $(x, t) \in \Pi(1)$  as  $\varepsilon \rightarrow 0$ .

*Proof.* Taking into account Lemma 1, it is sufficient to prove that  $u^\varepsilon - w^\varepsilon \rightarrow 0$  pointwise for  $(x, t) \in \Pi(1)$  as  $\varepsilon \rightarrow 0$ . Fix an arbitrary  $(x_0, t_0) \in \Pi(1)$ . If  $\varepsilon$  is sufficiently small, the point  $(x_0, t_0)$  belongs to  $\Pi^\varepsilon(1)$ , where we have the following integral representation:

$$\begin{aligned} (u^\varepsilon - w^\varepsilon)(x, t) &= c_r(\theta(x, t)) \left[ \int_0^L b_s^\varepsilon(\xi)(u^\varepsilon - w^\varepsilon)(\xi, \tau) d\xi \right. \\ &+ \left. \int_0^{Q(\tau)} b_r(\xi)(u^\varepsilon - w^\varepsilon)(\xi, \tau) d\xi + \int_{Q(\tau)}^L b_r(\xi)(u^\varepsilon - w^\varepsilon - v^\varepsilon)(\xi, \tau) d\xi \right] \Big|_{\tau=\theta(x, t)} \\ &+ \int_{\theta(x, t)}^t (u^\varepsilon - w^\varepsilon)(\xi, \tau) \left[ p(\xi, \tau) + \int_0^1 (\nabla_u f)(\xi, \tau, \sigma u^\varepsilon + (1 - \sigma)w^\varepsilon) d\sigma \right] \Big|_{\xi=\omega(\tau; x, t)} d\tau. \end{aligned} \quad (36)$$

Here

$$Q(t) = \begin{cases} \omega(t; 0, 0) & \text{if } \theta(L, t) > 0, \\ L & \text{if } \theta(L, t) = 0. \end{cases} \quad (37)$$

Note that

$$\omega(t; 0, \tau) \leq (t - \tau) \max_{(x,t) \in \Pi^T} \lambda(x, t). \quad (38)$$

Since, for small enough  $\varepsilon$ ,

$$|(u^\varepsilon - w^\varepsilon)(x_0, t_0)| \leq \max_{(x,t) \in \Pi^\varepsilon(1)} |(u^\varepsilon - w^\varepsilon)(x, t)|,$$

it is sufficient to prove that

$$\max_{(x,t) \in \Pi^\varepsilon(1)} |(u^\varepsilon - w^\varepsilon)(x, t)| = O(\varepsilon).$$

We start from the evaluation of the third integral in (36). On the account of (31), we represent it in the form

$$\begin{aligned} \int_{Q(t)}^L b_r(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx &= \int_{[Q(t), L] \times \{t\} \setminus (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})} b_r(u^\varepsilon - w^\varepsilon) dx \\ &+ \int_{[Q(t), L] \times \{t\} \cap (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})} b_r(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx. \end{aligned} \quad (39)$$

If  $\theta(L, t) = 0$ , this integral is equal to 0. Consider the case that  $\theta(L, t) > 0$ . To estimate the difference  $u^\varepsilon - w^\varepsilon$  on  $[Q(t), L] \times \{t\} \setminus (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})$ , we consider the corresponding problem

$$\begin{aligned} (\partial_t + \lambda(x, t)\partial_x)(u^\varepsilon - w^\varepsilon) &= \left(p(x, t) + \int_0^1 (\nabla_u f)(x, t, \sigma u^\varepsilon + (1 - \sigma)w^\varepsilon) d\sigma\right)(u^\varepsilon - w^\varepsilon), \\ (u^\varepsilon - w^\varepsilon)|_{t=0} &= 0. \end{aligned} \quad (40)$$

By Assumption 7 this problem has only the trivial solution. Therefore the second integral in (39) is equal to 0. We now estimate the third integral. We have the integral equation

$$\begin{aligned} &(u^\varepsilon - v^\varepsilon - w^\varepsilon)(x, t) \\ &= \int_0^t (p + F)(\xi, \tau)(u^\varepsilon - w^\varepsilon - v^\varepsilon)(\xi, \tau)|_{\xi=\omega(\tau, x, t)} d\tau \\ &+ \int_0^t [f(\xi, \tau, u^\varepsilon) - f(\xi, \tau, u^\varepsilon - v^\varepsilon)]|_{\xi=\omega(\tau, x, t)} d\tau \end{aligned}$$

that corresponds to the Cauchy problem (33)–(34). Combining it with Assumption 7, we conclude that

$$\max_{(x,t) \in \Pi_0^T \cap \overline{I_+^\varepsilon}} |u^\varepsilon - v^\varepsilon - w^\varepsilon| \leq C_1 \quad (41)$$

for a positive constant  $C_1$  not depending on  $\varepsilon$ . Therefore the absolute value of the third integral in (39) is bounded from above by  $C_1\varepsilon$ . As a consequence,

$$\left| \int_{Q(t)}^L b_r(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx \right| \leq C_2\varepsilon. \quad (42)$$

In the rest of the proof,  $C_i$  for  $i \geq 1$  are positive constants that do not depend on  $\varepsilon$ . We will distinguish two cases.

*Case 1.*  $\text{supp } b_s^{\varepsilon_0} \subset [\omega(t_1^*; 0, 0), L]$ . The first integral in (36) vanishes, since

$$\int_0^L b_s^\varepsilon(u^\varepsilon - w^\varepsilon) dx = \int_{[Q(t), L] \times \{t\} \setminus (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})} b_s^\varepsilon(u^\varepsilon - w^\varepsilon) dx \quad (43)$$

and  $u^\varepsilon - w^\varepsilon$  on  $[Q(t), L] \times \{t\} \setminus (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})$  satisfies the problem (40) which has only the trivial solution. Taking into account (7), (36)–(38), and (42), similarly to [13, p. 644] we obtain the following estimate that holds on  $\overline{\Pi^\varepsilon(1)} \cap \overline{\Pi^{t^0}}$ :

$$|(u^\varepsilon - w^\varepsilon)(x, t)| \leq \frac{C_2\varepsilon}{1 - q^0 t^0},$$

where

$$\begin{aligned} q^0 &= \max_{(x, t, y) \in \overline{\Pi^T} \times \mathbb{R}} |(\nabla_u f)(x, t, y)| \\ &+ \max_{(x, t) \in \overline{\Pi^T}} |p(x, t)| + \max_{t \in [0, T]} |c_r(t)| \max_{x \in [0, L]} |b_r(x)| \max_{(x, t) \in \overline{\Pi^T}} |\lambda(x, t)|, \\ t^0 &< q^0. \end{aligned}$$

Iterating this estimate at most  $\lceil T/t^0 \rceil$  times, each time using the final estimate for  $|u^\varepsilon - w^\varepsilon|$  from a preceding iteration, we obtain the following bound that holds on  $\overline{\Pi^\varepsilon(1)}$ :

$$|(u^\varepsilon - w^\varepsilon)(x, t)| \leq C_3\varepsilon. \quad (44)$$

This completes the proof in Case 1.

*Case 2.*  $\text{supp } b_s^{\varepsilon_0} \not\subset [\omega(t_1^*; 0, 0), L]$ . We fix an arbitrary sequence  $0 = p_0 < p_1 < p_2 < \dots < p_M = t_1^*$  such that  $\text{supp } b_s^{\varepsilon_0} \subset [\omega(p_j; 0, p_{j-1}), L]$ . Since  $\text{supp } b_s^\varepsilon \subset \text{supp } b_s^{\varepsilon_0}$  for  $\varepsilon \leq \varepsilon_0$ , we can choose the same sequence for all  $\varepsilon \leq \varepsilon_0$ . Given this sequence, we divide  $\overline{\Pi^\varepsilon(1)}$  into a finite number of subsets

$$\Pi^\varepsilon(1, j) = \left\{ (x, t) \in \overline{\Pi^\varepsilon(1)} \mid \tilde{\omega}(x; 0, p_{j-1}) \leq t \leq \tilde{\omega}(x; 0, p_j) \right\}. \quad (45)$$

We prove (44) with an appropriate choice of  $C_3$  separately for each of  $\Pi^\varepsilon(1, j)$ . Since  $\text{supp } b_s^\varepsilon \subset [\omega(p_1; 0, 0), L]$ , the conditions of Case 1 are true for  $\Pi^\varepsilon(1, 1)$ , and therefore the estimate (44) is true for this subset. The analog of (44) for  $\Pi^\varepsilon(1, 2)$  can be obtained in much the same way. We concentrate only on changes. For the first

integral in (36) we use the representation (43) with one more summand in the right hand side

$$\int_{\omega(t;0,p_1)}^{Q(t)} b_s^\varepsilon(u^\varepsilon - w^\varepsilon) dx.$$

The absolute value of this integral is bounded from above by  $C_4\varepsilon$  due to (44) on  $\Pi^\varepsilon(1,1)$ .

To derive (42) on  $\Pi^\varepsilon(1,2)$  with  $\omega(t;0,p_1)$  in place of  $Q(t)$  and with new  $C_2$ , we observe that in the analog of (39) there appears the third summand

$$\int_{\omega(t;0,p_1)}^{Q(t)} b_r(u^\varepsilon - w^\varepsilon) dx$$

that can be bounded from above by using (44) for  $\Pi^\varepsilon(1,1)$ . Similar arguments apply to all subsequent  $\Pi^\varepsilon(1,j)$ . Thus the estimate (44) is true for the whole  $\Pi^\varepsilon(1)$ .

The proof of Claim 2 is complete.  $\square$

**Claim 3** *The functions  $u^\varepsilon - w^\varepsilon - v^\varepsilon$  are bounded on  $\overline{I_+^\varepsilon(1)}$ , uniformly in  $\varepsilon$ .*

*Proof.* Two cases are possible.

*Case 1.*  $(0, t_1) \in I_+(1)$ . We have

$$(u^\varepsilon - v^\varepsilon - w^\varepsilon)|_{x=0} = G^\varepsilon(t) + (c_s^\varepsilon + c_r) \int_0^{\omega(t;0,t_1-\varepsilon)} b_r(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx, \quad (46)$$

$$t \in [t_1 - \varepsilon, t_1 + \varepsilon],$$

where

$$\begin{aligned} G^\varepsilon(t) = & (c_s^\varepsilon + c_r) \int_{\omega(t;0,t_1-\varepsilon)}^{Q(t)} (b_s^\varepsilon + b_r)(u^\varepsilon - w^\varepsilon) dx + (c_s^\varepsilon + c_r) \int_{Q(t)}^L b_s^\varepsilon(u^\varepsilon - w^\varepsilon) dx \\ & + (c_s^\varepsilon + c_r) \int_{Q(t)}^L b_r(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx. \end{aligned} \quad (47)$$

This representation follows from (32), (25), and (26). We now show that  $G^\varepsilon(t)$  is bounded on  $t \in [t_1 - \varepsilon, t_1 + \varepsilon]$ . Since  $[\omega(t;0,t_1 - \varepsilon), Q(t)] \times \{t\} \subset \overline{\Pi^\varepsilon(1)}$  for  $t \in [t_1 - \varepsilon, t_1 + \varepsilon]$ , the estimate (44) on  $\overline{\Pi^\varepsilon(1)}$  applies for the difference  $u^\varepsilon - w^\varepsilon$  under the first integral in (47). Using also (7) and (8), we conclude that the first summand in (47) is bounded uniformly in  $\varepsilon$ . Since  $u^\varepsilon - w^\varepsilon \equiv 0$  on  $\Pi_0^T \setminus I_+^\varepsilon$  (see (40)), the second integral is equal to 0. Applying (42) and (7) to the third summand, we see that  $G^\varepsilon(t)$  are bounded on  $[t_1 - \varepsilon, t_1 + \varepsilon]$ , uniformly in  $\varepsilon$ .

Observe that

$$\begin{aligned}
(u^\varepsilon - v^\varepsilon - w^\varepsilon)(x, t) &= (u^\varepsilon - v^\varepsilon - w^\varepsilon)(0, \theta(x, t)) \\
&+ \int_{\theta(x, t)}^t p(\xi, \tau)(u^\varepsilon - v^\varepsilon - w^\varepsilon)(\xi, \tau)|_{\xi=\omega(\tau; x, t)} d\tau \\
&+ \int_{\theta(x, t)}^t [f(\xi, \tau, u^\varepsilon) - f(\xi, \tau, w^\varepsilon)]|_{\xi=\omega(\tau; x, t)} d\tau, \quad (x, t) \in \overline{I_+^\varepsilon(1)},
\end{aligned} \tag{48}$$

where the boundary function  $(u^\varepsilon - w^\varepsilon - v^\varepsilon)(0, t)$  is given by (46). By Gronwall's argument applied to  $|(u^\varepsilon - v^\varepsilon - w^\varepsilon)(x, t)|$ , we easily obtain the estimate

$$\begin{aligned}
|(u^\varepsilon - v^\varepsilon - w^\varepsilon)(x, t)| &\leq C_3 \left[ 2T \max_{(x, t, y) \in \Pi^T \times \mathbb{R}} |f(x, t, y)| + \max_{(x, t) \in \overline{I_+^\varepsilon(1)}} |G^\varepsilon(\tau)| \right. \\
&\left. + (c_s^\varepsilon + c_r)(\tau) \int_0^{\omega(\tau; 0, t_1 - \varepsilon)} b_r(\xi)(u^\varepsilon - v^\varepsilon - w^\varepsilon)(\xi, \tau) d\xi \right]_{\tau=\theta(x, t)}, \quad (x, t) \in \overline{I_+^\varepsilon(1)}.
\end{aligned} \tag{49}$$

By (38) we have

$$\omega(t; 0, t_1^* - \varepsilon_1^-(\varepsilon)) \leq C_5 \varepsilon \tag{50}$$

for  $t \in [t_1^* - \varepsilon_1^-(\varepsilon), t_1^* + \varepsilon_1^+(\varepsilon)]$ . Given a mollifier  $\varphi(t)$ , let

$$q(\varepsilon) = C_3 C_5 \max_{x \in [0, \omega(t_1^* + \varepsilon_1^+(\varepsilon); 0, t_1^* - \varepsilon_1^-(\varepsilon))]} |b_r(x)| \left( \max_{t \in [0, T]} |\varphi(t)| + \varepsilon \max_{t \in [0, T]} |c_r(t)| \right). \tag{51}$$

By Assumptions 3 and 5,

$$\lim_{\varepsilon \rightarrow 0} q(\varepsilon) = 0. \tag{52}$$

We choose  $\varepsilon$  so small that

$$q(\varepsilon) < 1. \tag{53}$$

On the account of (49), (51), and (53), for sufficiently small  $\varepsilon$  we obtain

$$\max_{(x, t) \in \overline{I_+^\varepsilon(1)}} |(u^\varepsilon - v^\varepsilon - w^\varepsilon)(x, t)| \leq \frac{C_3}{1 - q(\varepsilon)} \left[ 2T \max_{(x, t, y) \in \Pi^T \times \mathbb{R}} |f(x, t, y)| + \max_{t \in [t_1 - \varepsilon, t_1 + \varepsilon]} |G^\varepsilon(t)| \right]. \tag{54}$$

*Case 2.*  $(0, t_1) \notin I_+(1)$ . By (32), (25), and (26) we have the equality (46) with  $t_1 - \varepsilon$  and  $t_1 + \varepsilon$  replaced by  $t_1^* - \varepsilon_1^-(\varepsilon)$  and  $t_1^* + \varepsilon_1^+(\varepsilon)$ , respectively, and with

$$\begin{aligned}
G^\varepsilon(t) &= c_r \int_{\omega(t; 0, t_1^* - \varepsilon_1^-(\varepsilon))}^{Q(t)} (b_s^\varepsilon + b_r)(u^\varepsilon - w^\varepsilon) dx + c_r \int_{Q(t)}^L b_s^\varepsilon (u^\varepsilon - w^\varepsilon) dx \\
&+ c_r \int_{Q(t)}^L b_r (u^\varepsilon - w^\varepsilon - v^\varepsilon) dx.
\end{aligned} \tag{55}$$

To estimate the absolute value of the first integral in (55) we apply (44) on  $\overline{\Pi^\varepsilon(1)}$  and (8). The second summand is equal to 0 (see (40)). For the third integral we use (42). It follows that  $G^\varepsilon(t)$  is bounded on  $[t_1^* - \varepsilon_1^-(\varepsilon), t_1^* + \varepsilon_1^+(\varepsilon)]$ , uniformly in  $\varepsilon$ . The rest of the proof runs as in Case 1, the minor changes being in using (49) and (54) with  $t_1^* - \varepsilon_1^-(\varepsilon)$  in place of  $t_1 - \varepsilon$  and  $t_1^* + \varepsilon_1^+(\varepsilon)$  in place of  $t_1 + \varepsilon$ .  $\square$

**Claim 4** 1. For every  $j \geq 1$ ,  $u^\varepsilon - w^\varepsilon - v^\varepsilon \rightarrow 0$  pointwise for  $(x, t) \in \Pi(j)$ , as  $\varepsilon \rightarrow 0$ .  
2. For every  $j \geq 1$ , the functions  $u^\varepsilon - w^\varepsilon - v^\varepsilon$  are bounded on  $\overline{I_+^\varepsilon(j)}$ , uniformly in  $\varepsilon$ .

*Proof.* Items 1 and 2 of the claim follow from the bounds

$$\max_{(x,t) \in \Pi^\varepsilon(j)} |(u^\varepsilon - w^\varepsilon)(x, t)| \leq A_j \varepsilon \quad (56)$$

and

$$\max_{(x,t) \in \overline{I_+^\varepsilon(j)}} |(u^\varepsilon - v^\varepsilon - w^\varepsilon)(x, t)| \leq B_j, \quad (57)$$

respectively, where  $A_j$  and  $B_j$  are constants depending only on  $j$ . We prove (56) and (57) by induction on  $j$ . The base case of  $j = 1$  is given by Claims 2 and 3. Assume that (56) and (57) are true for all  $j < i$ ,  $i \geq 2$ , and prove these estimates for  $j = i$ .

To prove (56) for  $j = i$ , we follow the proof of Claim 2 with the following changes. We use the formula (36) with  $\omega(\tau; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon))$  in place of  $Q(\tau)$ . To estimate the third integral in the analog of (36), we represent it in the form

$$\int_{\omega(t; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon))}^{Q(t)} b_r(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx + \int_{Q(t)}^L b_r(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx,$$

and apply the induction assumptions and (42). As a consequence, we obtain the estimate (42) with  $Q(t)$  replaced by  $\omega(t; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon))$  and with new  $C_2$ . Similarly to Claim 2, we distinguish two cases.

*Case 1.*  $\text{supp } b_s^{\varepsilon_0} \subset [\omega(t_i^*; 0, t_{i-1}^* - \varepsilon_{i-1}^-(\varepsilon_0)), L]$ . On the account of (40), we can rewrite the first summand in the analog of (36) in the form

$$\int_0^L b_s^\varepsilon(u^\varepsilon - w^\varepsilon) dx = \int_{[\omega(t_i^* - \varepsilon_i^-(\varepsilon); 0, t_{i-1}^* - \varepsilon_{i-1}^-(\varepsilon)), Q(t)] \times \{t\} \setminus (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})} b_s^\varepsilon(u^\varepsilon - w^\varepsilon) dx.$$

Applying (7) and (56) for  $j < i$ , we conclude that the absolute value of the integral is bounded from above by  $C_7 \varepsilon$ . The rest of the proof for this case runs similarly to the proof of Claim 2 in Case 1.

*Case 2.*  $\text{supp } b_s^{\varepsilon_0} \not\subset [\omega(t_i^*; 0, t_{i-1}^* - \varepsilon_{i-1}^-(\varepsilon_0)), L]$ . We fix an arbitrary sequence  $t_{i-1}^* = p_0 < p_1 < p_2 < \dots < p_M = t_i^*$  such that  $\text{supp } b_s^{\varepsilon_0} \subset [\omega(p_j; 0, p_{j-1}), L]$ . Given this sequence, we divide  $\overline{\Pi^\varepsilon(i)}$  into a finite number of subsets

$$\Pi^\varepsilon(i, j) = \{(x, t) \in \overline{\Pi^\varepsilon(i)} \mid \tilde{\omega}(x; 0, p_{j-1}) \leq t \leq \tilde{\omega}(x; 0, p_j)\}. \quad (58)$$

Observe that the partition of  $\overline{\Pi^\varepsilon(i)}$  is finite for every  $\varepsilon > 0$  and the number of subsets does not depend on  $\varepsilon$ . We further apply arguments similar to those used in the proof of Claim 2 for Case 2.

To prove (57) for  $j = i$ , we follow the proof of Claim 3 with the following changes. Similarly to Claim 3, we distinguish two cases.



*Case 1.*  $(0, t_k) \in I_+(i)$  for some  $k \leq l$ . We use the formula (46) with  $t_k$  in place of  $t_1$  and with

$$\begin{aligned} G^\varepsilon(t) = & (c_s^\varepsilon + c_r) \int_{[\omega(t; 0, t_i^* - \varepsilon_i^-(\varepsilon)), L] \times \{t\} \setminus (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})} (b_s^\varepsilon + b_r)(u^\varepsilon - w^\varepsilon) dx \\ & + (c_s^\varepsilon + c_r) \int_{[\omega(t; 0, t_i^* - \varepsilon_i^-(\varepsilon)), L] \times \{t\} \cap (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})} b_r(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx. \end{aligned} \quad (59)$$

Estimation of the first summand is based on the inclusion

$$[\omega(t; 0, t_i^* - \varepsilon_i^-(\varepsilon)), L] \times \{t\} \setminus (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\}) \subset \bigcup_{j=1}^i \overline{\Pi^\varepsilon(j)} \cup (\overline{\Pi_0^T} \setminus I_+^\varepsilon)$$

and (56), which for  $j < i$  is given by the induction assumptions and for  $j = i$  is just proved. Estimation of the second summand is based on the inclusion

$$[\omega(t; 0, t_i^* - \varepsilon_i^-(\varepsilon)), L] \times \{t\} \cap (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\}) \subset \bigcup_{j=1}^{i-1} \overline{I_+^\varepsilon(j)} \cup (\overline{\Pi_0^T} \cap \overline{I_+^\varepsilon}),$$

$$t \in [t_1 - \varepsilon, t_1 + \varepsilon],$$

and (57) for  $j < i$ .

*Case 2.*  $(0, t_k) \notin I_+(i)$  for all  $k \leq l$ . We use the formula (46) with  $t_i^* - \varepsilon_i^-(\varepsilon)$  and  $t_i^* + \varepsilon_i^+(\varepsilon)$  in place of  $t_1 - \varepsilon$  and  $t_1 + \varepsilon$ , respectively, and with

$$\begin{aligned} G^\varepsilon(t) = & c_r \int_{[\omega(t; 0, t_i^* - \varepsilon_i^-(\varepsilon)), L] \times \{t\} \setminus (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})} (b_s^\varepsilon + b_r)(u^\varepsilon - w^\varepsilon) dx \\ & + c_r \int_{[\omega(t; 0, t_i^* - \varepsilon_i^-(\varepsilon)), L] \times \{t\} \cap (I_+^\varepsilon \cap \{(x, t) \mid x \in \mathbb{R}\})} (b_s^\varepsilon + b_r)(u^\varepsilon - w^\varepsilon - v^\varepsilon) dx. \end{aligned} \quad (60)$$

In order to prove the boundedness of  $G^\varepsilon(t)$  we apply (56) for  $j \leq i$  and (57) for  $j < i$ .

The rest of the proof for both cases runs similarly to the proof of Claim 4 in Case 1.  $\square$

From Claim 4, (56) for  $j \leq n(T)$ , and (31) we conclude that the family  $(u^\varepsilon - w^\varepsilon - v^\varepsilon)_{\varepsilon > 0}$  is bounded on  $\overline{\Pi_1^T}$  uniformly in  $\varepsilon$  and converges to 0 almost everywhere in  $\overline{\Pi_1^T}$ . By dominated convergence theorem this family converges to 0 in  $L^1(\Pi_1^T)$ -norm. On the account of Claim 1,  $(u^\varepsilon - w^\varepsilon - v^\varepsilon)_{\varepsilon > 0}$  converges to 0 in  $L^1(\Pi^T)$ -norm. Since  $T$  is arbitrary, this is precisely the assertion of Lemma 2.

## 6 Proof of Lemma 3

Given  $T > 0$ , we choose  $\varepsilon_0$  so small that, for all  $\varepsilon \leq \varepsilon_0$ , the conditions (32) and

$$q(\varepsilon) \exp\left\{T \max_{(x, t) \in \Pi^T} |p(x, t)|\right\} < 1 \quad (61)$$

are fulfilled. Here  $q(\varepsilon)$  is defined by (51). The condition (61) follows from (52).

**Claim 1** *The family of functions  $w^\varepsilon$  converges in  $C(\overline{\Pi_0^T})$  as  $\varepsilon \rightarrow 0$ .*

*Proof.* For  $w^\varepsilon$  on  $\overline{\Pi_0^T}$  we use the representation given by (18) and (19). Since  $(Rw^\varepsilon)(x, t) = a_r(\omega(0; x, t))$  on  $\overline{\Pi_0^T}$ , the function  $w^\varepsilon$  for each  $\varepsilon > 0$  satisfies the same Volterra integral equation of the second kind. This means that  $w^\varepsilon$  does not depend on  $\varepsilon$  and for each  $\varepsilon > 0$  is equal to the same continuous function  $w(x, t)$  that can be found from the integral equation (18) by the method of sequential approximation. The claim follows.  $\square$

Therewith we are done in  $\overline{\Pi_0^T}$ . Since  $\overline{\Pi^T} = \overline{\Pi_0^T} \cup \overline{\Pi_1^T}$  and  $w^\varepsilon$  for each  $\varepsilon > 0$  is continuous on  $\overline{\Pi^T}$  (see Proposition 1 in Section 2), it remains to prove the convergence of  $w^\varepsilon$  in  $C(\overline{\Pi_1^T})$ . We will check the Cauchy criterion of the uniform convergence of  $w^\varepsilon$ . Given  $\delta > 0$ , we have to show for some  $\varepsilon_2 = \varepsilon_2(\delta)$  and every  $\varepsilon_1 < \varepsilon_2$  that

$$\left| (w^{\varepsilon_1} - w^{\varepsilon_2})(x, t) \right| \leq \delta \quad (62)$$

for all  $(x, t) \in \overline{\Pi_1^T}$ .

Because of so strong interaction of the regular and the singular parts (see the problems (11)–(13) and (14)–(16)), in the course of the proof of (62) we will need in parallel to prove some properties of  $v^\varepsilon$ .

Let

$$D_k(\varepsilon) = \int_{t_k^* - \varepsilon_k^-(\varepsilon)}^{t_k^* + \varepsilon_k^+(\varepsilon)} |v^\varepsilon(0, t)| dt \quad (63)$$

and

$$R_k(\varepsilon_1, \varepsilon_2) = \int_{t_k^* - \varepsilon_k^-(\varepsilon_2)}^{t_k^* + \varepsilon_k^+(\varepsilon_2)} (v^{\varepsilon_1} - v^{\varepsilon_2})(0, t) dt. \quad (64)$$

We will prove by induction on  $j$  the following 5 assertions for  $1 \leq j \leq n(T)$  ( $n(T)$  as well as  $\Pi(k)$  below are defined by Definition 3). Recall that  $n(T)$  does not depend on  $\varepsilon_2$ . Throughout this section  $C$  is a large enough constant that does not depend on  $\varepsilon$ .

**Assertion 1.** For every  $\delta > 0$ , if  $\varepsilon_2$  is small enough and  $\varepsilon_1 < \varepsilon_2$ , then (62) is true for all  $\overline{\Pi^{\varepsilon_2}(j)} \cup \overline{I_+^{\varepsilon_2}(j)}$ .

**Assertion 2.** The functions  $w^\varepsilon$  are bounded on  $\overline{\Pi^\varepsilon(j)} \cup \overline{I_+^\varepsilon(j)}$ , uniformly in  $\varepsilon > 0$ .

**Assertion 3.** The estimate  $D_j(\varepsilon) \leq C$  is true for all  $\varepsilon > 0$ .

**Assertion 4.** If  $\varepsilon_2$  is small enough and  $\varepsilon_1 < \varepsilon_2$ , then  $|R_j(\varepsilon_1, \varepsilon_2)| \leq C\varepsilon_2$ .

**Assertion 5.**  $w^\varepsilon(x, t)$  converges in  $C(\bigcup_{k=1}^j \overline{\Pi(k)} \cup \overline{\Pi_0^T})$  as  $\varepsilon \rightarrow 0$ .

Assertion 1 implies the Cauchy criterion of the uniform convergence of  $w^\varepsilon$  on  $\overline{\Pi_1^T}$ . Indeed, given  $\delta > 0$ , let  $\varepsilon_2$  be so small that (62) is true for every  $\varepsilon_1 < \varepsilon_2$  on

each  $\overline{\Pi^{\varepsilon_2}(j)} \cup \overline{I_+^{\varepsilon_2}(j)}$  for  $j \leq n(T)$ . Recall that, for any  $\varepsilon_2 > 0$ ,

$$\bigcup_{j=1}^{n(T)} (\overline{\Pi^{\varepsilon_2}(j)} \cup \overline{I_+^{\varepsilon_2}(j)}) = \overline{\Pi_1^T}.$$

It follows that (62) is true on  $\overline{\Pi_1^T}$  for all  $\varepsilon_1 < \varepsilon_2$ . By the Cauchy criterion,  $w^\varepsilon$  uniformly converges on  $\overline{\Pi_1^T}$ .

The proof of Assertions 1–5 for  $j = 1$  will be given by Claims 2–10. The induction step will be carried out by Claims 11–19.

To prove Assertion 1, we split  $\overline{\Pi^{\varepsilon_2}(j)} \cup \overline{I_+^{\varepsilon_2}(j)}$  into four subsets:

$$\overline{\Pi^{\varepsilon_2}(j)} \cup \overline{I_+^{\varepsilon_2}(j)} = \overline{\Pi^{\varepsilon_2}(j)} \cup (\overline{\Pi^{\varepsilon_1}(j)} \cap \overline{I_+^{\varepsilon_2}(j)}) \cup \overline{I_+^{\varepsilon_1}(j)} \cup (\overline{I_+^{\varepsilon_2}(j)} \cap \overline{\Pi^{\varepsilon_1}(j+1)}),$$

where each two neighboring subsets have common border. We will prove Assertion 1 separately for each of the four subsets.

**Claim 2** *The functions  $w^\varepsilon(x, t)$  are bounded on  $\overline{\Pi^\varepsilon(1)}$ , uniformly in  $\varepsilon$ .*

*Proof.* We use the representation of  $w^\varepsilon$  by (17) and (19) restricted to  $\overline{\Pi^\varepsilon(1)}$ . In this representation, on the account of (13), (21), and (22), we have

$$\begin{aligned} (Rw^\varepsilon)(x, t) &= c_r(\theta(x, t)) \int_0^{\omega(\theta(x, t); 0, 0)} (b_s^\varepsilon + b_r)(\xi) w^\varepsilon(\xi, \theta(x, t)) d\xi \\ &+ c_r(\theta(x, t)) \int_{\omega(\theta(x, t); 0, 0)}^L \left[ (b_s^\varepsilon + b_r)(\xi) w(\xi, \tau) + S(\xi, \tau) a_s^\varepsilon(\omega(0; \xi, \tau)) \right] \Big|_{\tau=\theta(x, t)} d\xi, \end{aligned}$$

where  $w(x, t) = w^\varepsilon(x, t)$  for all  $(x, t) \in \overline{\Pi_0^T}$  and for all  $\varepsilon > 0$  (see the proof of Claim 1). Taking into account (38) and the fact that  $\theta(x, t) \leq t$ , similarly to [13, p. 646] we obtain the global estimate

$$\begin{aligned} \max_{(x, t) \in \overline{\Pi^\varepsilon(1)}} |w^\varepsilon(x, t)| &\leq \left( \frac{1}{1 - q^1 t^1} \right)^{\lceil \frac{T}{t^1} \rceil} P(E) \left( 1 + \max_{x \in [0, L]} |a_r(x)| \right. \\ &\quad \left. + \max_{(x, t) \in \overline{\Pi^T}} |S(x, t)| \max_{\varepsilon} \int_0^L |a_s^\varepsilon(x)| dx \right), \end{aligned} \tag{65}$$

where

$$\begin{aligned} E &= \max_{t \in [0, T]} |c_r(t)| \left( \max_{x \in [0, L]} |b_r(x)| + \max_{\varepsilon} \int_0^L |b_s^\varepsilon(x)| dx \right), \\ q^1 &= (1 + LE) \left( \max_{(x, t, y) \in \overline{\Pi^T} \times \mathbb{R}} |(\nabla_u f)(x, t, y)| + \max_{(x, t) \in \overline{\Pi^T}} |p(x, t)| \right) + E \max_{(x, t) \in \overline{\Pi^T}} |\lambda(x, t)|, \end{aligned} \tag{66}$$

$t^1$  is a real so small that

$$t^1 < q^1 \tag{67}$$

and

$$\text{supp } b_s^\varepsilon \subset (\omega(kt^1; 0, (k-1)t^1), L] \quad \text{for all } 1 \leq k \leq \lceil T/t^1 \rceil, \quad (68)$$

and  $P(E)$  is a polynomial of degree  $\lceil T/t^1 \rceil$  with positive coefficients depending on  $f(x, t, 0)$ ,  $L$ , and  $T$ . The claim now follows by (8) and Assumptions 5 and 7.  $\square$

**Claim 3** 1. *Provided  $\varepsilon_2$  is small enough, for all  $\varepsilon'_2 \leq \varepsilon_2$  and for all  $\varepsilon'_1 \leq \varepsilon'_2$  the estimate*

$$\left| (w^{\varepsilon'_1} - w^{\varepsilon'_2})(x, t) \right| \leq \delta \quad (69)$$

*is true on  $\overline{\Pi^{\varepsilon_2}(1)}$ .*

2. *Provided  $\varepsilon_2$  is small enough, (62) is true on  $\overline{\Pi^{\varepsilon_2}(1)}$ .*

*Proof.* Recall that  $\overline{\Pi^{\varepsilon_2}(1)} \subset \overline{\Pi^{\varepsilon'_1}(1)}$  and  $\overline{\Pi^{\varepsilon_2}(1)} \subset \overline{\Pi^{\varepsilon'_2}(1)}$ . To represent  $w^{\varepsilon'_1}$  and  $w^{\varepsilon'_2}$  on  $\overline{\Pi^{\varepsilon_2}(1)}$ , we will use the system (11)–(13) restricted to  $\overline{\Pi^{\varepsilon_2}(1)}$ . For the difference  $w^{\varepsilon'_1} - w^{\varepsilon'_2}$  we will employ the corresponding linearized integral-operator equation. We distinguish two cases.

*Case 1.*  $\text{supp } b_s^{\varepsilon_2} \subset [\omega(t_1^*; 0, 0), L]$ . Using the fact that  $w^\varepsilon(x, t) \equiv w(x, t)$  on  $\overline{\Pi_0^T}$  for  $\varepsilon > 0$ , we obtain the integral equation

$$\begin{aligned} (w^{\varepsilon'_1} - w^{\varepsilon'_2})(x, t) = c_r(\theta(x, t)) \int_0^{\omega(\theta(x, t); 0, 0)} b_r(\xi) (w^{\varepsilon'_1} - w^{\varepsilon'_2})(\xi, \theta(x, t)) d\xi \\ + S_1(x, t) + S_2(\theta(x, t)) + S_3(\theta(x, t)), \end{aligned} \quad (70)$$

where

$$\begin{aligned} S_1(x, t) &= \int_{\theta(x, t)}^t \left[ p(\xi, \tau) \right. \\ &\quad \left. + \int_0^1 (\nabla_u f)(\xi, \tau, \sigma w^{\varepsilon'_1} + (1 - \sigma)w^{\varepsilon'_2}) d\sigma \right] (w^{\varepsilon'_1} - w^{\varepsilon'_2})(\xi, \tau) \Big|_{\xi=\omega(\tau; x, t)} d\tau, \\ S_2(t) &= c_r(t) \int_{Q(t)}^L (b_s^{\varepsilon'_1} - b_s^{\varepsilon'_2})(x) w(x, t) dx, \\ S_3(t) &= c_r(t) \int_{Q(t)}^L b_r(x) (v^{\varepsilon'_1} - v^{\varepsilon'_2})(x, t) dx, \end{aligned} \quad (71)$$

and  $Q(t)$  is defined by (37). We now estimate  $|S_2(t)|$  and  $|S_3(t)|$ . Since the function  $w$  on  $\overline{\Pi_0^T}$  is uniformly continuous, the properties (10) and (8) hold, and  $\text{supp } b_s^\varepsilon \subset \bigcup_{j=1}^k [x_j - \varepsilon, x_j + \varepsilon]$ , we have

$$\begin{aligned} |S_2(t)| &\leq \max_{t \in [0, T]} |c_r(t)| \int_{Q(t)}^L \left( |b_s^{\varepsilon'_1}| + |b_s^{\varepsilon'_2}| \right) \sum_{j=1}^k \left| w(x, t) - \chi_{[x_j - \varepsilon'_2, x_j + \varepsilon'_2]}(x) w(x_j, t) \right| dx \\ &\quad + \max_{t \in [0, T]} |c_r(t)| \sum_{j=1}^k \max_{(x_j, t) \in \overline{\Pi_0^T}} |w(x_j, t)| \left| \int_{x_j - \varepsilon'_2}^{x_j + \varepsilon'_2} (b_s^{\varepsilon'_1} - b_s^{\varepsilon'_2})(x) dx \right| \leq C\varepsilon_2. \end{aligned} \quad (72)$$

Here  $\chi_\Omega(x, t)$  denotes the characteristic function of a set  $\Omega$ .

Taking into account (22) and changing coordinates  $(x, t)$  to  $(\omega(0; x, t), t)$ , we estimate  $|S_3(t)|$  in the following way:

$$\begin{aligned}
|S_3(t)| &= \left| c_r(t) \int_{Q(t)}^L (b_r S)(x, t) (a_s^{\varepsilon'_1} - a_s^{\varepsilon'_2}) (\omega(0; x, t)) dx \right| \\
&= \left| c_r(t) \int_0^{\omega(0; L, t)} \frac{(b_r S)(\xi, t)}{(\partial_x \omega)(0; \xi, t)} \Big|_{\xi=\omega(t; x, 0)} (a_s^{\varepsilon'_1} - a_s^{\varepsilon'_2})(x) dx \right| \\
&\leq \max_{t \in [0, T]} |c_r(t)| \int_0^{x_{m(t)}^* + \varepsilon'_2} (|a_s^{\varepsilon'_1}| + |a_s^{\varepsilon'_2}|) \\
&\quad \times \sum_{j=1}^{m(t)} \left| \frac{(b_r S)(\omega(t; x, 0), t)}{(\partial_x \omega)(0; \omega(t; x, 0), t)} - \chi_{[x_j^* - \varepsilon'_2, x_j^* + \varepsilon'_2]}(x) \frac{(b_r S)(\omega(t; x_j^*, 0), t)}{(\partial_x \omega)(0; \omega(t; x_j^*, 0), t)} \right| dx \\
&+ \max_{t \in [0, T]} |c_r(t)| \sum_{j=1}^{m(t)} \max_{(\omega(t; x_j^*, 0), t) \in \overline{\Pi_0^0}} \left| \frac{(b_r S)(\omega(t; x_j^*, 0), t)}{(\partial_x \omega)(0, \omega(t; x_j^*, 0), t)} \right| \int_0^{x_{m(t)}^* + \varepsilon'_2} (a_s^{\varepsilon'_1} - a_s^{\varepsilon'_2})(x) dx \\
&\quad + \max_{t \in [0, T]} |c_r(t)| \int_{x_{m(t)}^* + \varepsilon'_2}^{\omega(0; L, t)} (|a_s^{\varepsilon'_1}| + |a_s^{\varepsilon'_2}|) \left| \frac{(b_r S)(\omega(t; x, 0), t)}{(\partial_x \omega)(0; \omega(t; x, 0), t)} \right| dx,
\end{aligned} \tag{73}$$

where  $m(t)$  is the number of indices  $j \leq m$  such that  $x_j^* + \varepsilon'_2 \in [0, \omega(0; L, t)]$ . Similarly to [13, p. 644] we obtain the estimate for  $|(w^{\varepsilon'_1} - w^{\varepsilon'_2})(x, t)|$  on  $\overline{\Pi^{\varepsilon_2}(1)} \cap \overline{\Pi^{t^1}}$ :

$$|(w^{\varepsilon'_1} - w^{\varepsilon'_2})(x, t)| \leq \frac{C\varepsilon_2}{1 - q^1 t^1}, \tag{74}$$

where  $q^1$  and  $t^1$  are defined by (66) and (67). Indeed, the second summand in the right-hand side of (73) is equal to 0 by (10). To estimate the first summand, we use (8) and the uniform continuity property for  $b_r$ ,  $S$ , and  $\lambda$  on  $\overline{\Pi^T}$ . To estimate the third summand, we observe that the integral is equal to 0 if  $\omega(0; L, t) \leq x_{m(t)}^* + \varepsilon'_2$  and is actually from  $x_{m(t)+1}^* - \varepsilon'_2$  to  $\omega(0; L, t)$ . In the latter case  $\omega(0; L, t) - x_{m(t)+1}^* + \varepsilon'_2 \leq C\varepsilon'_2$ . Combining this bound with the continuity of  $\lambda$  and the condition  $\omega(t; \omega(0; L, t), 0) = L$ , we obtain  $L - \omega(t; x_{m(t)+1}^* - \varepsilon'_2, 0) \leq C\varepsilon'_2$ . Since  $b_r(L) = 0$  by Assumption 3, we conclude that

$$\begin{aligned}
\max_{x \in [x_{m(t)+1}^* - \varepsilon'_2, \omega(0; L, t)]} |(b_r S)(\omega(t; x, 0), t)| &= \max_{x \in [\omega(t; x_{m(t)+1}^* - \varepsilon'_2, 0), L]} |(b_r S)(x, t)| \\
&\leq \max_{(x, t) \in [L - C\varepsilon'_2, L] \times [0, T]} |(b_r S)(x, t)| \leq C\varepsilon_2.
\end{aligned}$$

It follows that

$$|S_3(t)| \leq C\varepsilon_2. \tag{75}$$

Using (70), (72), and (75), we derive (74) by Gronwall's argument applied to  $|w^{\varepsilon'_1} - w^{\varepsilon'_2}|$ . Iterating this estimate at most  $\lceil T/t^1 \rceil$  times, each time using the final

estimate for  $|(w^{\varepsilon'_1} - w^{\varepsilon'_2})(x, t)|$  from a preceding iteration, we obtain the bound

$$\left| (w^{\varepsilon'_1} - w^{\varepsilon'_2})(x, t) \right| \leq \left( \frac{1}{1 - q^1 t^1} \right)^{\lceil \frac{T}{t^1} \rceil} P(E_1) C \varepsilon_2, \quad (76)$$

where

$$E_1 = \max_{(x, t) \in \Pi^T} |c_r(t) b_r(x)|$$

and  $P(E_1)$  is a polynomial of degree  $\lceil T/t^1 \rceil$  with positive coefficients depending on  $L$  and  $T$ .

*Case 2.*  $\text{supp } b_s^{\varepsilon_2} \not\subset [\omega(t_1^*; 0, 0), L]$ . We divide  $\overline{\Pi^{\varepsilon_2}(1)}$  into a finite number of subsets  $\Pi^{\varepsilon_2}(1, j)$ ,  $j \leq M$ , defined by (45) with  $\varepsilon$  replaced by  $\varepsilon_2$ . Note that, if  $j < M$ , then  $\Pi^{\varepsilon_2}(1, j)$  actually does not depend on  $\varepsilon_2$ . We prove an analog of (76) with an appropriate choice of  $t^1$ ,  $P$ , and  $C$ , separately for each of  $\Pi^{\varepsilon_2}(1, j)$ . Since  $\text{supp } b_s^{\varepsilon_2} \subset [\omega(p_1; 0, 0), L]$ , the conditions of Case 1 are true for  $\Pi^{\varepsilon_2}(1, 1)$ , and therefore the estimate (76) is true for this subset. Thus, provided  $\varepsilon_2$  is small enough, the estimate (69) holds on  $\Pi^{\varepsilon_2}(1, 1)$ .

The analog of (76) for  $\Pi^{\varepsilon_2}(1, 2)$  can be obtained in much the same way. We concentrate only on changes. On  $\Pi^{\varepsilon_2}(1, 2)$  we use the integral equation

$$\begin{aligned} (w^{\varepsilon'_1} - w^{\varepsilon'_2})(x, t) &= c_r(\theta(x, t)) \int_0^{\omega(\theta(x, t); 0, p_1)} b_r(\xi) (w^{\varepsilon'_1} - w^{\varepsilon'_2})(\xi, \theta(x, t)) d\xi \\ &+ S_1(x, t) + S_2(\theta(x, t)) + S_3(\theta(x, t)) + S_4(\theta(x, t)), \end{aligned} \quad (77)$$

where

$$S_4(t) = c_r(t) \int_{\omega(t; 0, p_1)}^{Q(t)} \left[ b_s^{\varepsilon'_1}(x) (w^{\varepsilon'_1} - w^{\varepsilon'_2})(x, t) + (b_s^{\varepsilon'_1} - b_s^{\varepsilon'_2})(x) w^{\varepsilon'_2}(x, t) \right] dx.$$

We now bound  $|S_2(t) + S_4(t)|$ . Observe that  $[\omega(t; 0, p_1), Q(t)] \times \{t\} \subset \Pi^{\varepsilon_2}(1, 1)$  if  $t \in [p_1, p_2]$ . By Proposition 1, Claim 1, and the estimate (69) on  $\Pi^{\varepsilon_2}(1, 1)$ , we conclude that  $w^\varepsilon$  converges in  $C(\Pi^{\varepsilon_2}(1, 1) \cup \overline{\Pi_0^T})$  to a continuous function  $w(x, t)$ . Using the equality  $w^{\varepsilon'_1}(x, t) = w^{\varepsilon'_2}(x, t) = w(x, t)$  on  $\overline{\Pi_0^T}$ , similarly to (72) we derive the bound

$$\begin{aligned} |S_2(t) + S_4(t)| &\leq \max_{t \in [0, T]} |c_r(t)| \max_{(x, t) \in \Pi(1, 1)} |w^{\varepsilon'_1} - w^{\varepsilon'_2}| \int_{\omega(t; 0, p_1)}^{Q(t)} |b_s^{\varepsilon'_1}| dx \\ &+ \max_{t \in [0, T]} |c_r(t)| \int_{\omega(t; 0, p_1)}^L \left( |b_s^{\varepsilon'_1}| + |b_s^{\varepsilon'_2}| \right) \left| w^{\varepsilon'_2}(x, t) - \sum_{j=1}^k \chi_{[x_j - \varepsilon'_2, x_j + \varepsilon'_2]}(x) w(x_j, t) \right| dx \\ &+ \max_{t \in [0, T]} |c_r(t)| \sum_{j=1}^k \max_{(x_j, t) \in \Pi_0^T \cup \Pi(1, 1)} |w(x_j, t)| \left| \int_{x_j - \varepsilon'_2}^{x_j + \varepsilon'_2} (b_s^{\varepsilon'_1} - b_s^{\varepsilon'_2})(x) dx \right| \leq C \varepsilon_2. \end{aligned}$$

Now, using (75), we conclude that (76) holds for  $\Pi^{\varepsilon_2}(1, 2)$  with new  $t^1$ ,  $P$ , and  $C$ .

Similar arguments apply to the subsets  $\Pi^{\varepsilon_2}(1, j)$ . Thus the estimate

$$\left| (w^{\varepsilon'_1} - w^{\varepsilon'_2})(x, t) \right| \leq C\varepsilon_2 \quad (78)$$

is true for the whole  $\overline{\Pi^{\varepsilon_2}(1)}$  in both cases.

The estimate (69) follows from (78), where  $\varepsilon_2$  is chosen small enough. The proof of Item 1 is complete.

Item 2 is a straightforward consequence of Item 1.  $\square$

**Claim 4** *Provided  $\varepsilon_2$  is small enough, (62) is true on  $\overline{\Pi^{\varepsilon_1}(1)} \cap \overline{I_+^{\varepsilon_2}(1)}$ .*

*Proof.* The functions  $w^{\varepsilon_1}(x, t)$  and  $w^{\varepsilon_2}(x, t)$  on  $\overline{\Pi^{\varepsilon_1}(1)} \cap \overline{I_+^{\varepsilon_2}(1)}$  are represented by (17) restricted, respectively, to  $\overline{\Pi^{\varepsilon_1}(1)}$  and  $\overline{I_+^{\varepsilon_2}(1)}$ .

Proposition 1 together with Claims 1 and 3 (Item 1) imply, for each fixed  $\varepsilon_2 > 0$  as small as in Claim 3, the Cauchy criterion of the uniform convergence of  $w^\varepsilon$  on  $\overline{\Pi_0^T} \cup \overline{\Pi^{\varepsilon_2}(1)}$ .

Therefore

$$w^\varepsilon(x, t) \text{ converges in } C(\overline{\Pi_0^T} \cup \overline{\Pi^{\varepsilon_2}(1)}) \text{ as } \varepsilon \rightarrow 0. \quad (79)$$

As above, the limit function will be denoted by  $w(x, t)$ . Now, using (32), we have the representation

$$\begin{aligned} & (w^{\varepsilon_1} - w^{\varepsilon_2})(x, t) \\ &= \left[ c_r(\tau) \int_{\omega(\tau; 0, t_1^* - \varepsilon_1^-(\varepsilon_2))}^L \left( (b_s^{\varepsilon_1} + b_r)(w^{\varepsilon_1} - w^{\varepsilon_2}) \right)(\xi, \tau) d\xi \right] \Big|_{\tau=\theta(x, t)} \\ & \quad + \left[ c_r(\tau) \int_{\omega(\tau; 0, t_1^* - \varepsilon_1^-(\varepsilon_2))}^L \left( b_s^{\varepsilon_1} - b_s^{\varepsilon_2} \right)(\xi) (w^{\varepsilon_2}(x, t) \right. \right. \\ & \quad \left. \left. - \sum_{j=1}^k \chi_{[x_j - \varepsilon_2, x_j + \varepsilon_2]}(x) w(x_j, t) \right) d\xi \right] \Big|_{\tau=\theta(x, t)} \\ & \quad + \left[ c_r(\tau) \sum_{j=1}^k w(x_j, t) \int_{\omega(\tau; 0, t_1^* - \varepsilon_1^-(\varepsilon_2))}^L \left( b_s^{\varepsilon_1} - b_s^{\varepsilon_2} \right)(\xi) d\xi \right] \Big|_{\tau=\theta(x, t)} + S_3(\theta(x, t)) \\ & \quad + \left[ c_r(\tau) \int_{\omega(\tau; 0, t_1^* - \varepsilon_1^-(\varepsilon_2))}^L \left( b_r(w^{\varepsilon_1} - w^{\varepsilon_2}) \right)(\xi, \tau) d\xi \right] \Big|_{\tau=\theta(x, t)} + S_1(x, t) + S_5(\theta(x, t)), \end{aligned} \quad (80)$$

where

$$S_5(t) = c_r(t) \int_0^{\omega(t; 0, t_1^* - \varepsilon_1^-(\varepsilon_2))} \left( b_r v^{\varepsilon_2} \right)(x, t) dx.$$

For the absolute values of the first four summands in (80) we obtain the upper bound  $C\varepsilon_2$  by the following argument. Note that  $[\omega(t; 0, t_1^* - \varepsilon_1^-(\varepsilon_2)), L] \times \{t\} \subset$

$\overline{\Pi_0^T} \cup \overline{\Pi^{\varepsilon_2}(1)}$  for  $t \in [t_1^* - \varepsilon_1^-(\varepsilon_2), t_1^* - \varepsilon_1^-(\varepsilon_1)]$ . For the first summand the bound now follows from (76), (8), and Claims 1 and 3 (Item 2). For the second summand we apply (79) and (8). For the third one we use the properties (10). For  $|S_3(t)|$  we use the estimate (75).

To prove the upper bound  $C\varepsilon_2$  for  $|S_5(t)|$  we need an estimate for  $v^\varepsilon$  on  $\overline{I_+^\varepsilon(1)}$ . To obtain it we consider two cases.

*Case 1.*  $(0, t_1) \in I_+(1)$ . Taking into account (32), we represent  $w^\varepsilon$  and  $v^\varepsilon$  on  $\overline{I_+^\varepsilon(1)}$  in the form

$$\begin{aligned} w^\varepsilon(x, t) &= [c_r(S_6^\varepsilon + S_7^\varepsilon)](\theta(x, t)) \\ &+ \int_{\theta(x, t)}^t f(\omega(\tau; x, t), \tau, w^\varepsilon) d\tau + \int_{\theta(x, t)}^t p(\omega(\tau; x, t), \tau) w^\varepsilon d\tau \end{aligned} \quad (81)$$

and

$$v^\varepsilon(x, t) = [c_s(S_6^\varepsilon + S_7^\varepsilon)](\theta(x, t)) + \int_{\theta(x, t)}^t p(\omega(\tau; x, t), \tau) v^\varepsilon d\tau, \quad (82)$$

where

$$\begin{aligned} S_6^\varepsilon(t) &= \int_{\omega(t; 0, t_1^* - \varepsilon_1^-(\varepsilon))}^L (b_s^\varepsilon + b_r)(x) w^\varepsilon(x, t) dx + \int_{Q(t)}^L (b_r v^\varepsilon)(x, t) dx, \\ S_7^\varepsilon(t) &= \int_0^{\omega(t; 0, t_1^* - \varepsilon_1^-(\varepsilon))} b_r(x) (w^\varepsilon + v^\varepsilon)(x, t) dx. \end{aligned} \quad (83)$$

Summing up, we have

$$\begin{aligned} w^\varepsilon(x, t) + v^\varepsilon(x, t) &= [(c_r + c_s^\varepsilon)S_6^\varepsilon](\theta(x, t)) + [(c_r + c_s^\varepsilon)S_7^\varepsilon](\theta(x, t)) \\ &+ \int_{\theta(x, t)}^t f(\omega(\tau; x, t), \tau, w^\varepsilon) d\tau + \int_{\theta(x, t)}^t p(\omega(\tau; x, t), \tau) (w^\varepsilon + v^\varepsilon) d\tau. \end{aligned} \quad (84)$$

By (21), (22), (8), Proposition 1, and Claim 2,  $S_6^\varepsilon(t)$  is a continuous function and satisfies the uniform in  $\varepsilon$  estimate

$$|S_6^\varepsilon(t)| \leq C, \quad t \in [t_1^* - \varepsilon_1^-(\varepsilon), t_1^* + \varepsilon_1^+(\varepsilon)]. \quad (85)$$

By Proposition 1,  $S_7^\varepsilon(t)$  is continuous. We now derive an upper bound for  $|S_7^\varepsilon(t)|$ . Applying the method of sequential approximation to the function  $w^\varepsilon + v^\varepsilon$  given by the formula (84), we obtain the estimate

$$\begin{aligned} \max_{(x, t) \in \overline{I_+^\varepsilon(1)}} |(w^\varepsilon + v^\varepsilon)(x, t)| &\leq \left[ T \max_{(x, t, y) \in \overline{\Pi^T} \times \mathbb{R}} |f(x, t, y)| \right. \\ &+ \max_{t \in [t_1 - \varepsilon, t_1 + \varepsilon]} |c_r(t) + c_s^\varepsilon(t)| \left( C + \max_{t \in [t_1 - \varepsilon, t_1 + \varepsilon]} |S_7^\varepsilon(t)| \right) \Big] \exp \left\{ T \max_{(x, t) \in \overline{\Pi^T}} |p(x, t)| \right\}. \end{aligned} \quad (86)$$



By (61),

$$\leq \frac{C}{1 - q(\varepsilon)C \exp\left\{T \max_{(x,t) \in \Pi^T} |p|\right\}} \left[ \max_{(x,t) \in \overline{I_+^\varepsilon(1)}} |w^\varepsilon + v^\varepsilon| + \max_{t \in [t_1 - \varepsilon, t_1 + \varepsilon]} |c_r(t) + c_s^\varepsilon(t)| \right]. \quad (87)$$

From (50), (87), (7), and Assumption 3 we conclude that

$$|S_7^\varepsilon(t)| \leq C \max_{x \in [0, \omega(t; 0, t_1 - \varepsilon)]} |b_r(x)| \leq C \max_{x \in [0, C_5 \varepsilon]} |b_r(x)| \leq C\varepsilon, \quad (88)$$

$$t \in [t_1^* - \varepsilon_1^-(\varepsilon), t_1^* + \varepsilon_1^+(\varepsilon)].$$

Combining (20) and (22) with (82), we obtain

$$v^\varepsilon(x, t) = c_s^\varepsilon(\theta(x, t)) (S_6^\varepsilon + S_7^\varepsilon)(\theta(x, t)) S(x, t), \quad (89)$$

where the functions  $|(S_6^\varepsilon + S_7^\varepsilon)(\theta(x, t))|$  on  $\overline{I_+^\varepsilon(1)}$  are bounded uniformly in  $\varepsilon$ . The latter is true by (85) and (88). The formula (89) implies

$$v^\varepsilon = O\left(\frac{1}{\varepsilon}\right) \quad (90)$$

for  $(x, t) \in \overline{I_+^\varepsilon(1)}$ . Taking into account (89), (50), (7), and Assumption 3, we derive the bound

$$|S_5(t)| \leq C \max_{x \in [0, \omega(t; 0, t_1 - \varepsilon_2)]} |b_r(x)| \leq C \max_{x \in [0, C_5 \varepsilon_2]} |b_r(x)| \leq C\varepsilon_2. \quad (91)$$

*Case 2.*  $(0, t_1) \notin I_+(1)$ . The proof is much the same as for Case 1. The only difference is in evaluation of  $|S_5(t)|$ , where  $v^\varepsilon$  on  $\overline{I_+^\varepsilon(1)}$  is now given by

$$v^\varepsilon(x, t) = \left[ c_r(\tau) \int_{\omega(\tau; 0, 0)}^L b_s^\varepsilon(\xi) S(\xi, \tau) a_s^\varepsilon(\omega(0; \xi, \tau)) d\xi \right] \Big|_{\tau=\theta(x, t)} S(x, t). \quad (92)$$

Hence (90) in this case is true by (7) and continuity of  $\lambda$ . Therefore for  $|S_5(t)|$  the estimate (91) holds.

We now return to (80) and, taking into account (91), estimate  $|w^{\varepsilon_1} - w^{\varepsilon_2}|$  following the proof of (76). As a result, the bound (62) is true for sufficiently small  $\varepsilon_2$ .  $\square$

**Claim 5** *Provided  $\varepsilon_2$  is small enough, (62) is true on  $\overline{I_+^{\varepsilon_1}(1)}$ .*

*Proof.* Note that  $w^{\varepsilon_1}$  and  $w^{\varepsilon_2}$  on  $\overline{I_+^{\varepsilon_1}(1)}$  are defined by the same formula (81). Therefore

$$(w^{\varepsilon_1} - w^{\varepsilon_2})(x, t) = [c_r(S_6^{\varepsilon_1} - S_6^{\varepsilon_2})](\theta(x, t)) + [c_r(S_7^{\varepsilon_1} - S_7^{\varepsilon_2})](\theta(x, t)) + S_1(x, t). \quad (93)$$

The upper bound  $C\varepsilon_2$  for the absolute value of the second summand follows from (88). To estimate the first summand, we use the equality

$$\begin{aligned} & (S_6^{\varepsilon_1} - S_6^{\varepsilon_2})(t) \\ &= \int_{\omega(t;0,t_1^*-\varepsilon_1^-(\varepsilon_2))}^L \left[ (b_s^{\varepsilon_1} + b_r)w^{\varepsilon_1} - (b_s^{\varepsilon_2} + b_r)w^{\varepsilon_2} + b_r(v^{\varepsilon_1} - v^{\varepsilon_2}) \right](x, t) dx \\ & \quad + \int_{\omega(t;0,t_1^*-\varepsilon_1^-(\varepsilon_1))}^{\omega(t;0,t_1^*-\varepsilon_1^-(\varepsilon_2))} (b_r w^{\varepsilon_1})(x, t) dx. \end{aligned}$$

The absolute value of the first summand is already estimated in the proof of Claim 4. For the second summand we can apply Claim 2, since  $[\omega(t;0,t_1^*-\varepsilon_1^-(\varepsilon_1)), \omega(t;0,t_1^*-\varepsilon_1^-(\varepsilon_2))] \times \{t\} \subset \overline{\Pi^{\varepsilon_1}(1)}$  for  $t \in [t_1^* - \varepsilon_1^-(\varepsilon_2), t_1^* + \varepsilon_1^+(\varepsilon_2)]$ . As a consequence,

$$|(S_6^{\varepsilon_1} - S_6^{\varepsilon_2})(t)| \leq C\varepsilon_2. \quad (94)$$

Applying the method of sequential approximation to  $w^{\varepsilon_1} - w^{\varepsilon_2}$  given by (93), on the account of (94) and the upper bound  $C\varepsilon_2$  for the second summand in (93), we derive the bound

$$|(w^{\varepsilon_1} - w^{\varepsilon_2})(x, t)| \leq C\varepsilon_2 \exp \left\{ T \left( \max_{(x,t,y) \in \overline{\Pi^T} \times \mathbb{R}} |(\nabla_u f)(x, t, y)| + \max_{(x,t) \in \overline{\Pi^T}} |p(x, t)| \right) \right\}. \quad (95)$$

This implies (62), provided  $\varepsilon_2$  is chosen small enough.  $\square$

**Claim 6** *Provided  $\varepsilon_2$  is small enough, (62) is true on  $\overline{I_+^{\varepsilon_2}(1)} \cap \overline{\Pi^{\varepsilon_1}(2)}$ .*

*Proof.* We follow the proof of Claim 4 with the following changes. On the account of (32), for  $w^{\varepsilon_1} - w^{\varepsilon_2}$  on  $\overline{I_+^{\varepsilon_2}(1)} \cap \overline{\Pi^{\varepsilon_1}(2)}$  we have the representation (80) with  $t_1^* - \varepsilon_1^-(\varepsilon_2)$  replaced by  $t_1^* + \varepsilon_1^+(\varepsilon_1)$  in the fifth summand, with

$$S_5(t) = c_r(t) \left[ \int_{\omega(t;0,t_1^*+\varepsilon_1^+(\varepsilon_1))}^{\omega(t;0,t_1^*-\varepsilon_1^-(\varepsilon_1))} (b_r v^{\varepsilon_1})(x, t) dx - \int_0^{\omega(t;0,t_1^*-\varepsilon_1^-(\varepsilon_2))} (b_r v^{\varepsilon_2})(x, t) dx \right], \quad (96)$$

and with one more summand

$$\left[ c_r(\tau) \int_{\omega(\tau;0,t_1^*+\varepsilon_1^+(\varepsilon_1))}^{\omega(\tau;0,t_1^*-\varepsilon_1^-(\varepsilon_2))} (b_r(w^{\varepsilon_1} - w^{\varepsilon_2}))(\xi, \tau) d\xi \right] \Big|_{\tau=\theta(x,t)}. \quad (97)$$

To estimate the absolute value of the latter expression, we use Claims 4 and 5. To estimate  $|S_5(t)|$ , to both integrals we apply the same argument that was used for evaluation of  $|S_5(t)|$  in the proof of Claim 4.  $\square$

By Claims 3–6, Assertion 1 is true for  $j = 1$ .

**Claim 7** *The functions  $w^\varepsilon$  are bounded on  $\overline{I_+^\varepsilon(1)}$ , uniformly in  $\varepsilon > 0$ .*

*Proof.* The claim follows from (81), (90), (50), and Assumptions 5 and 7.  $\square$

By Claims 2 and 7, Assertion 2 is true for  $j = 1$ .

**Claim 8** *The family of functions  $w^\varepsilon$  converges in  $C(\overline{\Pi_0^T} \cup \overline{\Pi(1)})$  as  $\varepsilon \rightarrow 0$ .*

*Proof.* Since, by Proposition 1, each  $w^\varepsilon$  is continuous, it suffices to prove the convergence separately on  $\overline{\Pi_0^T}$  and  $\overline{\Pi(1)}$ . On the former domain the convergence is given by Claim 2. The convergence on the latter domain follows by the Cauchy criterion which holds by Claims 3–5, and the fact that  $\overline{\Pi(1)} \subset \overline{\Pi^{\varepsilon_2}(1)} \cup \overline{I_+^{\varepsilon_2}(1)}$  for every  $\varepsilon_2 > 0$ . Thus, Assertion 5 is true for  $j = 1$ .

In the sequel the limit function will be denoted by  $w(x, t)$ .  $\square$

**Claim 9** *The estimate  $D_1(\varepsilon) \leq C$  is true for all  $\varepsilon > 0$ .*

*Proof.* *Case 1.*  $(0, \tilde{t}_1) \in I_+(1)$ . Then

$$\begin{aligned} D_1(\varepsilon) &= \int_{t_1^* - \varepsilon_1^-(\varepsilon)}^{t_1^* + \varepsilon_1^+(\varepsilon)} \left| c_r(t) \int_{Q(t)}^L b_s^\varepsilon(x) S(x, t) a_s^\varepsilon(\omega(0; x, t)) dx \right| dt \\ &\leq \max_{(x, t) \in \overline{\Pi_0^T}} \left| \frac{(c_r S)(x, t)}{(\partial_t \omega)(0; x, t)} \right| \int_0^L |a_s^\varepsilon(x)| dx \int_0^L |b_s^\varepsilon(x)| dx \leq C. \end{aligned}$$

The estimate follows from (8) and Assumptions 5 and 6.

*Case 2.*  $(0, t_1) \in I_+(1)$ . Then

$$D_1(\varepsilon) = \int_{t_1 - \varepsilon}^{t_1 + \varepsilon} |c_s^\varepsilon(t)(S_6^\varepsilon + S_7^\varepsilon)(t)| dt \leq C.$$

This estimate follows from (85), (88), and (8).

Thus, Assertion 3 is true for  $j = 1$ .  $\square$

**Claim 10** *If  $\varepsilon_2$  is small enough and  $\varepsilon_1 < \varepsilon_2$ , then*

$$|R_1(\varepsilon_1, \varepsilon_2)| \leq C\varepsilon_2.$$

*Proof.* We will consider  $\varepsilon_2$  as small as in Claims 3–6. We will use (16) restricted to  $[t_1^* - \varepsilon_1^-(\varepsilon_2), t_1^* + \varepsilon_1^+(\varepsilon_2)]$  and (22).

*Case 1.*  $(0, \tilde{t}_1) \in I_+(1)$ . By (25) and (26) we have  $c_s^\varepsilon = 0$ , and therefore

$$R_1(\varepsilon_1, \varepsilon_2) = \int_{t_1^* - \varepsilon_1^-(\varepsilon_2)}^{t_1^* + \varepsilon_1^+(\varepsilon_2)} c_r(t) \int_{Q(t)}^L (b_s^{\varepsilon_1} v^{\varepsilon_1} - b_s^{\varepsilon_2} v^{\varepsilon_2})(x, t) dx dt.$$

Applying (22) restricted to  $\overline{\Pi_0^T}$  and changing coordinates  $(x, t) \rightarrow (x, \xi) = (x, \omega(0; x, t))$ , we obtain

$$R_1(\varepsilon_1, \varepsilon_2) = \sum_{(q,d) \in E} \int_{x_d^* - \varepsilon_2}^{x_d^* + \varepsilon_2} \int_{x_q - \varepsilon_2}^{x_q + \varepsilon_2} Q(x, \xi) [b_s^{\varepsilon_1}(x) a_s^{\varepsilon_1}(\xi) - b_s^{\varepsilon_2}(x) a_s^{\varepsilon_2}(\xi)] dx d\xi,$$

where

$$Q(x, \xi) = \frac{(c_r S)(x, t)}{(\partial_t \omega)(0; x, t)} \Big|_{t=\tilde{\omega}(x; \xi, 0)}$$

is a continuous function in  $x$  and  $\xi$  and  $E$  is the set of pairs of indices  $q \leq k$  and  $d \leq m$  such that  $\omega(0; x_q, \tilde{t}_1) = x_d^*$ . Evidently, if  $(0, \tilde{t}_1) \in I_+(1)$ , then there exists at least one pair  $(q, d)$  that satisfies the latter condition. Then

$$\begin{aligned} |R_1(\varepsilon_1, \varepsilon_2)| &\leq \sum_{q,d} \max_{(x,\xi) \in [x_q - \varepsilon_2, x_q + \varepsilon_2] \times [x_d^* - \varepsilon_2, x_d^* + \varepsilon_2]} |Q(x, \xi) - Q(x_q, x_d^*)| \\ &\quad \times \left[ \int_{x_q - \varepsilon_2}^{x_q + \varepsilon_2} |b_s^{\varepsilon_1}| dx \int_{x_d^* - \varepsilon_2}^{x_d^* + \varepsilon_2} |a_s^{\varepsilon_1}| dx + \int_{x_q - \varepsilon_2}^{x_q + \varepsilon_2} |b_s^{\varepsilon_2}| dx \int_{x_d^* - \varepsilon_2}^{x_d^* + \varepsilon_2} |a_s^{\varepsilon_2}| dx \right] \\ &\quad + \sum_{q,d} |Q(x_q, x_d^*)| \left| \int_{x_q - \varepsilon_2}^{x_q + \varepsilon_2} b_s^{\varepsilon_1} dx \int_{x_d^* - \varepsilon_2}^{x_d^* + \varepsilon_2} (a_s^{\varepsilon_1} - a_s^{\varepsilon_2}) dx + \int_{x_d^* - \varepsilon_2}^{x_d^* + \varepsilon_2} a_s^{\varepsilon_2} dx \int_{x_q - \varepsilon_2}^{x_q + \varepsilon_2} (b_s^{\varepsilon_1} - b_s^{\varepsilon_2}) dx \right|. \end{aligned}$$

The last two summands are equal to 0 by (10). The first summand is bounded from above by  $C\varepsilon_2$  by (8) and the continuity property for  $Q(x, t)$ . This completes the proof in Case 1.

*Case 2.*  $(0, t_1) \in I_+(1)$ . Then the second summand in (16) is equal to 0, and therefore

$$R_1(\varepsilon_1, \varepsilon_2) = \int_{t_1 - \varepsilon_2}^{t_1 + \varepsilon_2} \left[ c_s^{\varepsilon_1}(t) (S_6^{\varepsilon_1} + S_7^{\varepsilon_1})(t) - c_s^{\varepsilon_2}(t) (S_6^{\varepsilon_2} + S_7^{\varepsilon_2})(t) \right] dt. \quad (98)$$

We hence have

$$\begin{aligned} |R_1(\varepsilon_1, \varepsilon_2)| &= \left| \int_{t_1 - \varepsilon_2}^{t_1 + \varepsilon_2} (c_s^{\varepsilon_1} S_7^{\varepsilon_1} - c_s^{\varepsilon_2} S_7^{\varepsilon_2})(t) dt + \int_{t_1 - \varepsilon_2}^{t_1 + \varepsilon_2} (c_s^{\varepsilon_1} - c_s^{\varepsilon_2})(t) (S_6^{\varepsilon_1}(t) - S_6^{\varepsilon_1}(t_1)) dt \right. \\ &\quad \left. + S_6^{\varepsilon_1}(t_1) \int_{t_1 - \varepsilon_2}^{t_1 + \varepsilon_2} (c_s^{\varepsilon_1} - c_s^{\varepsilon_2})(t) dt + \int_{t_1 - \varepsilon_2}^{t_1 + \varepsilon_2} c_s^{\varepsilon_2}(t) (S_6^{\varepsilon_1} - S_6^{\varepsilon_2})(t) dt \right| \\ &\leq \int_{t_1 - \varepsilon_2}^{t_1 + \varepsilon_2} (|c_s^{\varepsilon_1}| |S_7^{\varepsilon_1}| + |c_s^{\varepsilon_2}| |S_7^{\varepsilon_2}|) dt + \int_{t_1 - \varepsilon_2}^{t_1 + \varepsilon_2} (|c_s^{\varepsilon_1}| + |c_s^{\varepsilon_2}|) |S_6^{\varepsilon_1}(t) - S_6^{\varepsilon_1}(t_1)| dt \\ &\quad + \int_{t_1 - \varepsilon_2}^{t_1 + \varepsilon_2} |c_s^{\varepsilon_2}| |S_6^{\varepsilon_1} - S_6^{\varepsilon_2})(t)| dt. \end{aligned} \quad (99)$$

Combining (88) with (8), we obtain the upper bound  $C\varepsilon_2$  for the first summand in the right-hand side of the inequality (99). The same bound for the last summand follows from (94) and (8). It remains to estimate the second summand. Without loss of generality we assume that  $t \geq t_1$ . By (32) we have the representation

$$\begin{aligned}
S_6^{\varepsilon_1}(t) - S_6^{\varepsilon_1}(t_1) = & \int_{\omega(t;0,t_1-\varepsilon_1)}^L (b_s^{\varepsilon_1} + b_r)(x) (w^{\varepsilon_1}(x, t) - w^{\varepsilon_1}(x, t_1)) dx \\
& + \int_{\omega(t_1;0,t_1-\varepsilon_1)}^{\omega(t;0,t_1-\varepsilon_1)} b_r(x) w^{\varepsilon_1}(x, t_1) dx + \int_{Q(t)}^L (b_r S)(x, t) a_s^{\varepsilon_1}(\omega(0; x, t)) dx \\
& - \int_{\omega(t_1;0,0)}^{\omega(t_1;L,t)} (b_r S)(x, t_1) a_s^{\varepsilon_1}(\omega(0; x, t_1)) dx - \int_{\omega(t_1;L,t)}^L (b_r S)(x, t_1) a_s^{\varepsilon_1}(\omega(0; x, t_1)) dx.
\end{aligned} \tag{100}$$

Changing coordinates  $x \rightarrow \omega(0; x, t)$  in the third integral and  $x \rightarrow \omega(0; x, t_1)$  in the fourth integral, after simple computation we have

$$\begin{aligned}
|S_6^{\varepsilon_1}(t) - S_6^{\varepsilon_1}(t_1)| \leq & \left( \int_0^L |b_s^{\varepsilon_1}| dx + L \max_{x \in [0,L]} |b_r(x)| \right) \\
& \times \max_{|t-t_1| \leq \varepsilon_1, (x,t) \in \overline{\Pi_0^T} \cup \overline{\Pi(1)}} \left[ |w^{\varepsilon_1} - w|(x, t)| + |w(x, t) - w(x, t_1)| \right. \\
& \left. + |w(x, t_1) - w^{\varepsilon_1}(x, t_1)| \right] \\
& + \max_{(x,t) \in \overline{\Pi^{\varepsilon_1}(1)}} |w^{\varepsilon_1}(x, t)| \max_{x \in [0,L]} |b_r(x)| |\omega(t; 0, t_1 - \varepsilon_1) - \omega(t_1; 0, t_1 - \varepsilon_1)| \\
& + \left[ \max_{(x,t) \in [0,L] \times [t_1-\varepsilon_1, t_1+\varepsilon_1]} \left| \frac{(b_r S)(\omega(t; x, 0), t)}{(\partial_x \omega)(0; \omega(t; x, 0), t)} - \frac{(b_r S)(\omega(t_1; x, 0), t_1)}{(\partial_x \omega)(0; \omega(t_1; x, 0), t_1)} \right| \right. \\
& \left. + \max_{x \in [\omega(t_1-\varepsilon_1; L, t_1+\varepsilon_1), L]} |b_r(x)| \max_{(x,t) \in \overline{\Pi^T}} |S(x, t)| \right] \int_0^L |a_s^{\varepsilon_1}| dx \leq C\varepsilon_2.
\end{aligned} \tag{101}$$

The latter estimate is true by Claims 2, 7, and 8, estimates (8), and Assumptions 3 and 5.

Claim 9 follows, and therefore Assertion 4 is true for  $j = 1$ .  $\square$

**Induction assumption.** We assume that Assertions 1–5 are true for  $j \leq i - 1$ ,  $i \geq 2$ .

**Claim 11** *The functions  $w^{\varepsilon}(x, t)$  are bounded on  $\overline{\Pi^{\varepsilon}(i)}$ , uniformly in  $\varepsilon > 0$ .*

*Proof.* The proof is similar to the proof of Claim 2. The function  $w^\varepsilon(x, t)$  on  $\overline{\Pi^\varepsilon(i)}$  is defined by the formula (17), where

$$\begin{aligned} (Rw^\varepsilon)(x, t) = & c_r(\theta(x, t)) \left[ \int_0^{\omega(\tau; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon))} (b_s^\varepsilon + b_r)(\xi) w^\varepsilon(\xi, \tau) d\xi \right. \\ & + \int_{\omega(\tau; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon))}^L (b_s^\varepsilon + b_r)(\xi) w^\varepsilon(\xi, \tau) d\xi + \int_{\omega(\tau; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon))}^{Q(\tau)} (b_r S)(\xi, \tau) v^\varepsilon(0, \theta(\xi, \tau)) d\xi \\ & \left. + \int_{Q(\tau)}^L (b_r S)(\xi, \tau) a_s^\varepsilon(\omega(0; \xi, \tau)) d\xi \right] \Big|_{\tau=\theta(x, t)}. \end{aligned}$$

Taking into account (8) and Assertions 2 and 3 for  $j < i$ , we conclude that the last three summands are bounded uniformly in  $\varepsilon$ . Similarly to [13, p. 646], we obtain the global estimate

$$\begin{aligned} & \max_{(x, t) \in \overline{\Pi_1^\varepsilon(i)}} |w^\varepsilon(x, t)| \\ & \leq \left( \frac{1}{1 - q^1 t^1} \right)^{\lceil \frac{T}{t^1} \rceil} P(E) \left( 1 + \max_{(x, t) \in \left( \bigcup_{j=1}^{i-1} \overline{\Pi(j)} \right) \cup I_+^{\varepsilon_1(i-1)} \cup \overline{\Pi_0^T}} |w^\varepsilon(x, t)| \right), \end{aligned} \quad (102)$$

where  $q^1$ ,  $t^1$ , and  $E$  are defined by (66), (67), and (68), and  $P(E)$  is a polynomial of degree  $\lceil T/t^1 \rceil$  with positive coefficients depending on  $f(x, t, 0)$ ,  $L$ , and  $T$ . The claim now follows from Assertion 2 for  $j \leq i - 1$ .  $\square$

**Claim 12** 1. *Provided  $\varepsilon_2$  is small enough, for all  $\varepsilon'_2 \leq \varepsilon_2$  and for all  $\varepsilon'_1 \leq \varepsilon'_2$  the estimate (69) is true on  $\overline{\Pi(i)} \setminus I_+^{\varepsilon_2(i)}$ .*  
2. *Provided  $\varepsilon_2$  is small enough, (62) is true on  $\overline{\Pi^{\varepsilon_2}(i)}$ .*

*Proof.* We fix an arbitrary sequence  $t_{i-1}^* = p_0 < p_1 < p_2 < \dots < p_M = t_i^* - \varepsilon_i^-(\varepsilon_2)$  such that  $M \geq 2$ ,  $p_1 > t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon_2)$ , and  $\text{supp } b_s^{\varepsilon_2} \subset [\omega(p_i; 0, p_{i-1}), L]$ . We can do so due to (32). Given this sequence, we divide  $\overline{\Pi(i)} \setminus I_+^{\varepsilon_2(i)}$  into a finite number of subsets

$$\Pi^{\varepsilon_2}(i, j) = \left\{ (x, t) \in \overline{\Pi(i)} \setminus I_+^{\varepsilon_2(i)} \mid \tilde{\omega}(x; 0, p_{j-1}) \leq t \leq \tilde{\omega}(x; 0, p_j) \right\}.$$

Note that, if  $j < M$ , then  $\Pi^{\varepsilon_2}(i, j)$  actually does not depend on  $\varepsilon_2$ .

To obtain (69) for  $\Pi^{\varepsilon_2}(i, 1)$ , we first derive (62) for  $\Pi^{\varepsilon_2}(i, 1)$ . Doing this, we follow the proof of Claim 3 (Item 1) for Case 1 with the following changes. Throughout the proof  $\varepsilon'_1$  and  $\varepsilon'_2$  are replaced by  $\varepsilon_1$  and  $\varepsilon_2$ , respectively. We use the formulas (70) and (71) with  $\omega(t; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon_2))$  in place of  $Q(t)$  and with

$$S_2(t) = c_r(t) \int_{\omega(t; 0, t_{i-1}^*)}^L \left[ (b_s^{\varepsilon_1} - b_s^{\varepsilon_2}) w^{\varepsilon_2} + b_s^{\varepsilon_1} (w^{\varepsilon_1} - w^{\varepsilon_2}) \right] (x, t) dx$$

$$+c_r(t) \int_{\omega(t;0,t_{i-1}^*+\varepsilon_{i-1}^+(\varepsilon_2))}^L b_r(x)(w^{\varepsilon_1}-w^{\varepsilon_2})(x,t) dx.$$

To estimate  $|S_2|$ , we use Assertions 1, 2, and 5 for  $j < i$ . To estimate the analog of  $|S_3|$ , we use Assertions 4 and 5 for  $j < i$ . As a result, (62) is true on  $\Pi^{\varepsilon_2}(i, 1)$ .

Note that  $\Pi^{\varepsilon_2}(i, 1)$  does not depend on  $\varepsilon_1$  and  $\varepsilon_2$ . By Assertion 1 for  $j = i - 1$  and (62) for  $\Pi^{\varepsilon_2}(i, 1)$ , that we have just proved, we conclude that  $w^\varepsilon$  converges in  $C(\Pi^{\varepsilon_2}(i, 1))$ . The estimate (69) is hence true on  $\Pi^{\varepsilon_2}(i, 1)$ .

The estimate (69) for  $\Pi^{\varepsilon_2}(i, 2)$  follows similarly to the proof of Claim 3 (Item 1) for Case 2 (see the proof of (69) for  $\Pi^{\varepsilon_2}(1, 2)$ ). The minor change is that we estimate  $|S_2(t) + S_4(t)|$  using Assertion 5 for  $j = i - 1$  and (69) for  $\Pi^{\varepsilon_2}(i, 1)$ , that we have just proved. Similar arguments apply to the subsets  $\Pi^{\varepsilon_2}(i, j)$ ,  $j \leq M$ . As a result, (69) is true for the whole  $\overline{\Pi(i)} \setminus I_+^{\varepsilon_2}(i)$ .

Item 1 follows.

Item 2 is straightforward consequence of Item 1.  $\square$

**Claim 13** *Provided  $\varepsilon_2$  is small enough, (62) is true on  $\overline{\Pi^{\varepsilon_1}(i)} \cap \overline{I_+^{\varepsilon_2}(i)}$ .*

*Proof.* The proof is similar to the proof of Claim 4. We concentrate only on the changes that here appear. We choose  $\varepsilon_2$  so small that the condition (32) with  $t_1^* + \varepsilon_1^+(\varepsilon_2)$  replaced by  $t_i^* + \varepsilon_i^+(\varepsilon_2)$  and  $t_1^* - \varepsilon_1^-(\varepsilon_2)$  replaced by  $t_i^* - \varepsilon_i^-(\varepsilon_2)$  is true. As in Claim 4, we distinguish two cases.

*Case 1.* *There exists  $j \leq l$  such that  $(0, t_j) \in I_+(i)$ .* We use (80) with  $t_j^*$  in place of  $t_1^*$ , where  $Q(t)$  in  $S_3(t)$  is replaced by  $\omega(t; 0, t_j - \varepsilon_2)$ . For the first four summands in the analog of (80) we use Assertions 1–4 for  $j < i$  and Claim 12. The upper bound for  $|S_5(t)|$  follows from the bounds

$$|S_6^\varepsilon(t)| \leq C, \quad t \in [t_i^* - \varepsilon_i^-(\varepsilon), t_i^* + \varepsilon_i^+(\varepsilon)] \quad (103)$$

and

$$|S_7^\varepsilon(t)| \leq C\varepsilon, \quad t \in [t_i^* - \varepsilon_i^-(\varepsilon), t_i^* + \varepsilon_i^+(\varepsilon)], \quad (104)$$

where  $S_6^\varepsilon(t)$  and  $S_7^\varepsilon(t)$  are defined by (83) with  $t_i^* - \varepsilon_i^-(\varepsilon)$  in place of  $t_1^* - \varepsilon_1^-(\varepsilon)$  and  $\omega(t; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon))$  in place of  $Q(t)$ . The estimate (103) is true by the arguments used for obtaining (85), Assertions 2 and 3 for  $j < i$ , and Claim 11. The estimate (104) is obtained similarly to (88). The estimates (103) and (104) imply (90).

*Case 2.*  $(0, t_j) \notin I_+(i)$  for  $j \leq l$ . On the account of (32), (25), and (26), on  $\overline{I_+^\varepsilon(i)}$  we have the equality

$$\begin{aligned} v^\varepsilon(x, t) = & \left[ c_r(\tau) \int_{\omega(\tau; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon_2))}^{Q(\tau)} b_s^\varepsilon(\xi) S(\xi, \tau) v^\varepsilon(0, \theta(\xi, \tau)) d\xi \right] \Big|_{\tau=\theta(x, t)} S(x, t) \\ & + \left[ c_r(\tau) \int_{Q(\tau)}^L b_s^\varepsilon(\xi) S(\xi, \tau) a_s^\varepsilon(\omega(0; \xi, \tau)) d\xi \right] \Big|_{\tau=\theta(x, t)} S(x, t). \end{aligned} \quad (105)$$

For the second summand we apply the same arguments as in the proof of Claim 4 for Case 2. For the first summand we apply (7), Assertion 3 for  $j < i$  and the inclusion

$$\begin{aligned} & \{(0, \theta(x, t)) \mid t \in [t_i - \varepsilon_i^-(\varepsilon), t_i + \varepsilon_i^+(\varepsilon)], x \in [\omega(t; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon_2)), Q(t)]\} \\ & \subset \bigcup_{j=1}^{i-1} (\overline{I_+^\varepsilon(j)} \cap \{(x, t) \mid x = 0\}). \end{aligned}$$

Therefore (90) is true on  $\overline{I_+^\varepsilon(i)}$ . The claim follows.  $\square$

From what has already been proved we conclude that (90) holds on  $\overline{I_+^\varepsilon(j)}$  for every  $j \leq n(T)$ . Since  $v^\varepsilon(x, t) = S(x, t)a_s^\varepsilon(\omega(0; x, t))$  for  $(x, t) \in \overline{\Pi_0^T}$ , (90) holds on  $\overline{\Pi_0^T}$ .

By (31),  $\text{supp } v^\varepsilon \subset \overline{I_+^\varepsilon}$ . Therefore

$$v^\varepsilon = O\left(\frac{1}{\varepsilon}\right) \quad \text{on} \quad \overline{\Pi^T}. \quad (106)$$

We will need this property in the sequel.

**Claim 14** *Provided  $\varepsilon_2$  is small enough, (62) is true on  $\overline{I_+^{\varepsilon_1}(i)}$ .*

*Proof.* We follow the proof of Claim 5 with the following changes. We use (83) and (93) with  $t_1^* - \varepsilon_1^-(\varepsilon)$  and  $Q(t)$  to be replaced by  $t_i^* - \varepsilon_i^-(\varepsilon)$ . To estimate the absolute value of the first summand in the analog of (93), we use Assertions 1–3 for  $j < i$  and Claims 11–13. As a consequence,

$$\left| (S_6^{\varepsilon_1} - S_6^{\varepsilon_2})(x, t) \right| \leq C\varepsilon_2, \quad t \in [t_i^* - \varepsilon_i^-(\varepsilon), t_i^* + \varepsilon_i^+(\varepsilon)]. \quad (107)$$

To estimate the second summand, we use (104).  $\square$

**Claim 15** *Provided  $\varepsilon_2$  is small enough, (62) is true on  $\overline{I_+^{\varepsilon_2}(i)} \cap \overline{\Pi^{\varepsilon_1}(i+1)}$ .*

*Proof.* We follow the proof of Claim 6 with the following changes. For  $w^{\varepsilon_1} - w^{\varepsilon_2}$  on  $\overline{I_+^{\varepsilon_2}(i)} \cap \overline{\Pi^{\varepsilon_1}(i+1)}$  we use (80) with  $t_1^* - \varepsilon_1^-(\varepsilon_2)$  replaced by  $t_i^* - \varepsilon_i^-(\varepsilon_2)$  in the fifth summand and with one more summand (97), where  $t_1^* + \varepsilon_1^+(\varepsilon_1)$  and  $t_1^* - \varepsilon_1^-(\varepsilon_2)$  are replaced by  $t_i^* + \varepsilon_i^+(\varepsilon_1)$  and  $t_i^* - \varepsilon_i^-(\varepsilon_2)$ , respectively. In the representation of  $S_3(t)$  we now have  $\omega(t; t_{i-1}^* - \varepsilon_{i-1}^-(\varepsilon_2))$  in place of  $Q(t)$ . For  $S_5(t)$  we now use the formula (96) with  $t_i^* - \varepsilon_i^-(\varepsilon_1)$ ,  $t_i^* + \varepsilon_i^+(\varepsilon_1)$ , and  $t_i^* - \varepsilon_i^-(\varepsilon_2)$  in place of  $t_1^* - \varepsilon_1^-(\varepsilon_1)$ ,  $t_1^* + \varepsilon_1^+(\varepsilon_1)$ , and  $t_1^* - \varepsilon_1^-(\varepsilon_2)$ , respectively.

To estimate the absolute value of the analog of (97) we use Claims 13 and 14.  $\square$

By Claims 12–15, Assertion 1 is true for  $j = i$ .

**Claim 16** *The functions  $w^\varepsilon(x, t)$  are bounded on  $\overline{I_+^\varepsilon(i)}$ , uniformly in  $\varepsilon > 0$ .*

*Proof.* Note that the function  $w^\varepsilon(x, t)$  on  $\overline{I_+^\varepsilon(i)}$  is defined by (81) and (83) with  $t_i^* - \varepsilon_i^-(\varepsilon)$  in place of  $t_1^* - \varepsilon_1^-(\varepsilon)$  and  $\omega(t; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon))$  in place of  $Q(t)$ . The claim follows from (103), (104), and Assumptions 5 and 7.  $\square$



Assertion 2 for  $j = i$  follows from Claims 11 and 16.

**Claim 17** *The family of functions  $w^\varepsilon$  converges in  $C(\bigcup_{j=1}^i \overline{\Pi(j)} \cup \overline{\Pi_0^T})$  as  $\varepsilon \rightarrow 0$ .*

*Proof.* Since, by Proposition 1, each  $w^\varepsilon$  is continuous, it suffices to prove the convergence separately on  $\overline{\Pi_0^T}$  and  $\bigcup_{j=1}^i \overline{\Pi(j)}$ . On the former domain the convergence is ensured by Claim 2. The convergence on the latter domain follows by the Cauchy criterion which holds by Assertion 1 for  $j \leq i$ , and the fact that  $\bigcup_{j=1}^i \overline{\Pi(j)} \subset \bigcup_{j=1}^i (\overline{\Pi^{\varepsilon_2}(j)} \cup I_+^{\varepsilon_2}(j))$  for every  $\varepsilon_2 > 0$ .  $\square$

By Claim 17, Assertion 5 holds for  $j = i$ .

**Claim 18** *The estimate  $D_i(\varepsilon) \leq C$  is true for all  $\varepsilon > 0$ .*

*Proof.* *Case 1.*  $(0, \tilde{t}_i) \in I_+(i)$ . We have

$$\begin{aligned} D_i(\varepsilon) &= \int_{t_i^* - \varepsilon_i^-(\varepsilon)}^{t_i^* + \varepsilon_i^+(\varepsilon)} \left| c_r(t) \int_0^{Q(t)} b_s^\varepsilon(x) S(x, t) v^\varepsilon(0, \theta(x, t)) dx \right. \\ &\quad \left. + c_r(t) \int_{Q(t)}^L b_s^\varepsilon(x) S(x, t) a_s^\varepsilon(\omega(0; x, t)) dx \right| dt \\ &\leq \max_{(x, t) \in \overline{\Pi_1^T}} \left| \frac{(c_r S)(x, t)}{(\partial_t \theta)(x, t)} \right| \sum_{j=1}^{i-1} \int_{t_j^* - \varepsilon_j^-(\varepsilon)}^{t_j^* + \varepsilon_j^+(\varepsilon)} |v^\varepsilon(0, t)| dt \int_0^L |b_s^\varepsilon(x)| dx \\ &\quad + \max_{(x, t) \in \overline{\Pi_0^T}} \left| \frac{(c_r S)(x, t)}{(\partial_t \omega)(0; x, t)} \right| \int_0^L |a_s^\varepsilon(x)| dx \int_0^L |b_s^\varepsilon(x)| dx \leq C, \end{aligned}$$

where  $Q(t)$  is defined by (37). This estimate is true by (8) and Assertion 3 for  $j < i$ .

*Case 2.*  $(0, t_j) \in I_+(i)$  for some  $j \leq l$ . We have

$$D_i(\varepsilon) = \int_{t_j - \varepsilon}^{t_j + \varepsilon} |c_s^\varepsilon(t)(S_6^\varepsilon + S_7^\varepsilon)(t)| dt \leq C,$$

This estimate is true by (103), (104), and (8).  $\square$

Claim 18 implies Assertion 3 for  $j = i$ .

**Claim 19** *The estimate  $|R_i(\varepsilon_1, \varepsilon_2)| \leq C\varepsilon_2$  is true for  $\varepsilon_2$  so small that Assertion 1 holds for all  $j \leq i$ .*

*Proof.* We follow the proof of Claim 10 with the changes listed below. Similarly to Claim 10, we distinguish two cases.

*Case 1.*  $(0, \tilde{t}_i) \in I_+(i)$ . We have  $R_i(\varepsilon_1, \varepsilon_2) = R_i^1(\varepsilon_1, \varepsilon_2) + R_i^2(\varepsilon_1, \varepsilon_2)$ , where

$$R_i^1(\varepsilon_1, \varepsilon_2) = \int_{t_i^* - \varepsilon_i^-(\varepsilon_2)}^{t_i^* + \varepsilon_i^+(\varepsilon_2)} c_r(t) \int_0^{Q(t)} (b_s^{\varepsilon_1} v^{\varepsilon_1} - b_s^{\varepsilon_2} v^{\varepsilon_2})(x, t) dx dt$$

and

$$R_i^2(\varepsilon_1, \varepsilon_2) = \int_{t_i^* - \varepsilon_i^-(\varepsilon_2)}^{t_i^* + \varepsilon_i^+(\varepsilon_2)} c_r(t) \int_{Q(t)}^L (b_s^{\varepsilon_1} v^{\varepsilon_1} - b_s^{\varepsilon_2} v^{\varepsilon_2})(x, t) dx dt.$$

If  $Q(t) < L$ , then  $[Q(t), L] \times \{t\} \subset \overline{\Pi_0^T}$ . Therefore  $|R_i^2(\varepsilon_1, \varepsilon_2)|$  can be estimated in the same way as  $|R_1(\varepsilon_1, \varepsilon_2)|$  was estimated in the proof of Claim 10 for Case 1. The minor change is that now  $E$  will denote the set of pairs of indices  $q \leq k$  and  $d \leq m$  such that  $\omega(0; x_q, \tilde{t}_i) = x_d^*$ . It remains to estimate  $|R_i^1(\varepsilon_1, \varepsilon_2)|$ . Applying (21) and (22) restricted to  $\overline{\Pi_1^T}$  and changing coordinates  $(x, t) \rightarrow (x, \xi) = (x, \theta(x, t))$ , we obtain

$$R_i^1(\varepsilon_1, \varepsilon_2) = \sum_{(q,d) \in J} \int_{t_d^* - \varepsilon_d^-(\varepsilon_2)}^{x_d + \varepsilon_d^+(\varepsilon_2)} \int_{x_q - \varepsilon_2}^{x_q + \varepsilon_2} Q_1(x, t) [(b_s^{\varepsilon_1} - b_s^{\varepsilon_2})(x) v^{\varepsilon_1}(0, t) - b_s^{\varepsilon_2}(x) (v^{\varepsilon_1} - v^{\varepsilon_2})(0, t)] dx dt,$$

where

$$Q_1(x, \xi) = \frac{(c_r S)(x, \tau)}{(\partial_t \theta)(x, \tau)} \Big|_{\tau = \tilde{\omega}(x, 0, t)}$$

and  $J$  is the set of pairs of indices  $q \leq k$  and  $d \leq i - 1$  such that  $\omega(x_q; 0, t_d^*) = \tilde{t}_i$ . Obviously, at least one of the sets  $E$  or  $J$  is nonempty. To estimate  $|R_i^1(\varepsilon_1, \varepsilon_2)|$ , in addition to the arguments used for estimation of  $|R_2(\varepsilon_1, \varepsilon_2)|$  we apply Assertions 3 and 4 for  $j < i$ .

*Case 2.*  $(0, t_j) \in I_+(i)$  for some  $j \leq l$ . We use (98) and (99) with  $t_1$  replaced by  $t_j$ . To estimate the first and the third summands in the analog of (99), we apply (8), (104), and (107). To estimate the second summand, we use the representation

$$\begin{aligned} S_6^{\varepsilon_1}(t) - S_6^{\varepsilon_1}(t_j) &= \int_{\omega(t; 0, t_j - \varepsilon_1)}^L (b_s^{\varepsilon_1} + b_r)(x) (w^{\varepsilon_1}(x, t) - w^{\varepsilon_1}(x, t_j)) dx \\ &\quad + \int_{\omega(t_j; 0, t_j - \varepsilon_1)}^{\omega(t; 0, t_j - \varepsilon_1)} b_r(x) w^{\varepsilon_1}(x, t_j) dx \\ &\quad + \left[ \int_{Q(t)}^L (b_r S)(x, t) a_s^{\varepsilon_1}(\omega(0; x, t)) dx - \int_{Q(t_j)}^L (b_r S)(x, t_j) a_s^{\varepsilon_1}(\omega(0; x, t_j)) dx \right] \end{aligned}$$

$$+ \left[ \int_{\omega(t;0,t_j-\varepsilon_1)}^{Q(t)} (b_r S)(x,t) v^{\varepsilon_1}(0, \theta(x,t)) dx - \int_{\omega(t_j;0,t_j-\varepsilon_1)}^{Q(t_j)} (b_r S)(x,t_j) v^{\varepsilon_1}(0, \theta(x,t_j)) dx \right]. \quad (108)$$

The absolute value of the first three summands are estimated similarly to estimation of  $|S_6^{\varepsilon_1}(t) - S_6^{\varepsilon_1}(t_1)|$  in the proof of Claim 10 (see (101)). In addition to the arguments used for (101), we apply Assertion 2 for  $j < i$  and Claims 11, 16, and 17. We now concentrate on the last summand. Let us rewrite it in the form

$$\begin{aligned} & \int_{\theta(Q(t),t)}^{t_j-\varepsilon_1} \left[ \frac{(b_r S)(x,t)}{(\partial_x \theta)(x,t)} \Big|_{x=\omega(t;0,\tau)} - \frac{(b_r S)(x,t_j)}{(\partial_x \theta)(x,t_j)} \Big|_{x=\omega(t_j;0,\tau)} \right] v^{\varepsilon_1}(0, \tau) d\tau \\ & + \int_{\theta(Q(t_j),t_j)}^{\theta(Q(t),t)} \frac{(b_r S)(x,t)}{(\partial_x \theta)(x,t)} \Big|_{x=\omega(t;0,\tau)} v^{\varepsilon_1}(0, \tau) d\tau. \end{aligned}$$

The absolute value of this expression is less than or equal to

$$\begin{aligned} & \left[ \max_{(x,t) \in [0,L] \times [t_j-\varepsilon_1, t_j+\varepsilon_1]} \left| \frac{(b_r S)(x,t)}{(\partial_x \theta)(x,t)} \Big|_{x=\omega(t;0,\tau)} - \frac{(b_r S)(x,t_j)}{(\partial_x \theta)(x,t_j)} \Big|_{x=\omega(t_j;0,\tau)} \right| \\ & + \max_{x \in [\omega(t_j-\varepsilon_1; Q(t_j+\varepsilon_1), t_j+\varepsilon_1), Q(t_j-\varepsilon_1)]} |b_r(x)| \max_{(x,t) \in \Pi^T} |S(x,t)| \Big] \\ & \times \sum_{r=1}^{i-1} \int_{t_r^* - \varepsilon_r^-(\varepsilon_1)}^{t_r^* + \varepsilon_r^+(\varepsilon_1)} |v^{\varepsilon_1}(0, t)| dt \leq C \varepsilon_2. \end{aligned}$$

The latter bound is true by Assertion 3 for  $j < i$  and Assumption 3.

We conclude that Assertion 4 holds for  $j = i$ .  $\square$

Thus the induction step is done and the proof of Lemma 3 is complete.

## 7 Proof of Lemma 4

Given an arbitrary test function  $\varphi \in \mathcal{D}(\Pi)$ , we have to show the convergence of  $\langle v^\varepsilon(x, t), \varphi(x, t) \rangle$  as  $\varepsilon \rightarrow 0$ . Fix  $T = T(\varphi) > 0$  such that  $\text{supp } \varphi \subset \Pi^T$ . Let

$$v^\varepsilon(x, t) = \sum_{i=0}^{\infty} v_i^\varepsilon(x, t), \quad (109)$$

where

$$\text{supp } v_0^\varepsilon(x, t) = \text{supp} \left\{ v^\varepsilon(x, t) \Big|_{\Pi_0 \cap I_+^\varepsilon} \right\}, \quad \Pi_0 = \{(x, t) \in \Pi \mid x > \omega(t; 0, 0)\},$$

and

$$\text{supp} \left\{ v_i^\varepsilon(x, t) \Big|_{\Pi^T} \right\} = \text{supp} \left\{ v^\varepsilon(x, t) \Big|_{I_+^\varepsilon(i)} \right\} \quad \text{for } i \leq \rho(T),$$

$\rho(T)$  is as in Definition 3. Clearly,  $v_i^\varepsilon(x, t)$  for  $i \geq 1$  is supported on one of the connected components of  $(\bar{\Pi} \setminus \Pi_0) \cap \bar{I}_+^\varepsilon$ . The representation (109) is true because  $\text{supp } v^\varepsilon \subset \bar{I}_+^\varepsilon$  by (31). Since  $\langle v^\varepsilon(x, t), \varphi(x, t) \rangle = \langle \sum_{i=0}^{\rho(T)} v_i^\varepsilon(x, t), \varphi(x, t) \rangle$ , it suffices to prove that  $\langle v_i^\varepsilon(x, t), \varphi(x, t) \rangle$  converges as  $\varepsilon \rightarrow 0$  separately for each  $0 \leq i \leq \rho(T)$ .

**Claim 1**  $\langle v_0^\varepsilon(x, t), \varphi(x, t) \rangle$  converges as  $\varepsilon \rightarrow 0$ .

*Proof.* By (21) and (22),

$$v_0^\varepsilon(x, t) = S(x, t) a_s^\varepsilon(\omega(0; x, t)). \quad (110)$$

Let us compute the action

$$\begin{aligned} \langle v_0^\varepsilon(x, t), \varphi(x, t) \rangle &= \int_{\Pi_0^T} S(x, t) a_s^\varepsilon(\omega(0; x, t)) \varphi(x, t) d(x, t) \\ &= \int_{\tilde{\Pi}_0^T} \frac{S(\xi, t) \varphi(\xi, t)}{(\partial_x \omega)(0; \xi, t)} \Big|_{\xi=\omega(t; x, 0)} a_s^\varepsilon(x) d(x, t), \end{aligned}$$

where

$$\tilde{\Pi}_0^T = \{(x, t) \in \mathbb{R}^2 \mid (\omega(t; x, 0), t) \in \Pi_0^T\}.$$

It follows easily that

$$\langle v_0^\varepsilon(x, t), \varphi(x, t) \rangle \xrightarrow{\varepsilon \rightarrow 0} \sum_{i=1}^m \int_{\tilde{\Pi}_0^T \cap \{(x, t) \mid x=x_i^*\}} \frac{S(\xi, t) \varphi(\xi, t)}{\omega_\xi(0; \xi, t)} \Big|_{\xi=\omega(t; x_i^*, 0)} dt,$$

Here we used a simple change of coordinates  $F_0 : (x, t) \rightarrow (\omega(0; x, t), t)$ , where  $F_0$  maps  $\Pi_0^T$  to  $\tilde{\Pi}_0^T$ . Since  $\tilde{\Pi}_0^T$  is a bounded domain, the claim is proved.  $\square$

**Claim 2**  $\langle v_1^\varepsilon(x, t), \varphi(x, t) \rangle$  converges as  $\varepsilon \rightarrow 0$ .

*Proof.* Two cases are possible.

*Case 1.*  $(0, t_1) \in I_+(1)$ . Let

$$S_8^\varepsilon(t) = \int_0^L [(b_s^\varepsilon + b_r) w^\varepsilon + b_r v^\varepsilon] dx.$$

From (13) we conclude that  $c_r(t) S_8^\varepsilon(t) = w^\varepsilon(0, t)$ . By Lemma 3, the family of functions  $w^\varepsilon(0, t)$  is uniformly convergent on  $[0, T]$ . Since  $c_r(t)$  is an arbitrary continuous function, the same assertion is true for the family of functions  $S_8^\varepsilon(t)$ . Hence there exists a continuous function  $S_8^0(t)$  such that

$$\lim_{\varepsilon \rightarrow 0} S_8^\varepsilon(t) = S_8^0(t) \text{ in } C[0, T].$$

Therefore, if  $|t - t_1| \leq C\varepsilon_2$ , we have

$$|S_8^\varepsilon(t) - S_8^0(t_1)| \leq |(S_8^\varepsilon - S_8^0)(t)| + |S_8^0(t) - S_8^0(t_1)| \leq C\varepsilon_2.$$

Note that  $S_8^\varepsilon(t) = S_6^\varepsilon(t) + S_7^\varepsilon(t)$  whenever  $t \in [t_1 - \varepsilon, t_1 + \varepsilon]$ . Using the representation (89), we conclude that

$$\begin{aligned} \langle v_1^\varepsilon(x, t), \varphi(x, t) \rangle &= \int_{I_+^{\varepsilon_0}(1)} c_s^\varepsilon(\theta(x, t)) S_8^\varepsilon(\theta(x, t)) S(x, t) \varphi(x, t) d(x, t) \\ &= \int_{\tilde{I}_+^{\varepsilon_0}(1)} c_s^\varepsilon(t) S_8^\varepsilon(t) \frac{S(x, \tau) \varphi(x, \tau)}{(\partial_t \tilde{\omega})(0; x, \tau)} \Big|_{\tau=\tilde{\omega}(x; 0, t)} d(x, t) \\ &\xrightarrow{\varepsilon \rightarrow 0} S_8^0(t_1) \int_{\tilde{I}_+^{\varepsilon_0}(1) \cap \{(x, t) \mid t=t_1\}} \frac{S(x, \tau) \varphi(x, \tau)}{(\partial_t \tilde{\omega})(0; x, \tau)} \Big|_{\tau=\tilde{\omega}(x; 0, t_1)} dx. \end{aligned}$$

Here

$$\tilde{I}_+^{\varepsilon_0}(1) = \{(x, t) \in \mathbb{R}^2 \mid (x, \tilde{\omega}(x; 0, t)) \in I_+^{\varepsilon_0}(1)\}$$

is a bounded domain. This implies that the latter integral is finite, and therefore the claim in this case is true.

*Case 2.*  $(0, t_1) \notin I_+(1)$ . Using the representation of  $v_1^\varepsilon$  given by (105), consider the action

$$\begin{aligned} &\langle v_1^\varepsilon(x, t), \varphi(x, t) \rangle \\ &= \langle c_r(\theta(x, t)) \int_{\omega(\theta(x, t); 0, 0)}^L b_s^\varepsilon(\xi) a_s^\varepsilon(\omega(0; \xi, \theta(x, t))) S(\xi, \theta(x, t)) d\xi S(x, t), \varphi(x, t) \rangle \\ &= \int_{I_+^{\varepsilon_0}(1)} \int_{\omega(\theta(x, t); 0, 0)}^L c_r(\theta(x, t)) b_s^\varepsilon(\xi) a_s^\varepsilon(\omega(0; \xi, \theta(x, t))) S(\xi, \theta(x, t)) S(x, t) \varphi(x, t) d\xi d(x, t). \end{aligned}$$

Changing coordinates  $(\xi, x, t) \rightarrow (\xi, x, \omega(0; \xi, \theta(x, t)))$ , we convert the latter expression into

$$\begin{aligned} &\int_{\Omega} b_s^\varepsilon(\xi) a_s^\varepsilon(t) \left[ c_r(\theta(x, \tau)) S(\xi, \theta(x, \tau)) S(x, \tau) \varphi(x, \tau) \right. \\ &\quad \left. \times [(\partial_t \omega)(0; \xi, \theta(x, \tau)) (\partial_t \theta)(x, \tau)]^{-1} \right] \Big|_{\tau=\tilde{\omega}(x; 0, \tilde{\omega}(\xi; t, 0))} d(\xi, x, t), \end{aligned}$$

where

$$\Omega = \{(\xi, x, t) \mid t = \omega(0; \xi, \theta(x, \tau)), (x, \tau) \in I_+^{\varepsilon_0}(1), \omega(\theta(x, \tau); 0, 0) < \xi < L\}$$

is a bounded domain,  $I(i)$  is the set of those pairs  $(j, r)$  that  $j \leq k$ ,  $r \leq m$ , and  $\omega(0; x_j, t_i^*) = x_r^*$ . It is now clear that, as  $\varepsilon \rightarrow 0$ ,  $\langle v_1^\varepsilon(x, t), \varphi(x, t) \rangle$  converges to

$$c_r(t_1^*) \sum_{(j, r) \in I(1)} \frac{S(x_j, t_1^*)}{(\partial_t \omega)(0; x_j, t_1^*)} \int_{\Omega \cap \{(\xi, x, t) \mid \xi=x_j, t=x_r^*\}} \frac{S(x, t) \varphi(x, t)}{(\partial_t \theta)(x, t)} \Big|_{t=\tilde{\omega}(x; 0, t_1^*)} dx.$$

Since the sum is finite, the claim follows.  $\square$

**Claim 3**  $\lim_{\varepsilon \rightarrow 0} v_1^\varepsilon(0, t) = C_1 \delta(t - t_1^*)$  in  $\mathcal{D}'(\mathbb{R}_+)$ , where  $C_1$  is a real constant.

*Proof.* Take a test function  $\psi(t) \in \mathcal{D}(\mathbb{R}_+)$  and compute the action  $\langle v_1^\varepsilon(0, t), \psi(t) \rangle$ . Similarly to the proof of Claim 2, we obtain that

$$v_1^\varepsilon(0, t) \xrightarrow{\varepsilon \rightarrow 0} S_8^0(t_1) S(0, t_1) \delta(t - t_1), \quad \text{if } (0, t_1) \in I_+(1),$$

and

$$v_1^\varepsilon(0, t) \xrightarrow{\varepsilon \rightarrow 0} c_r(t_1^*) \sum_{\{j \leq k \mid (j, r) \in I(1), r \leq m\}} \frac{S(x_j, t_1^*)}{(\partial_t \omega)(0; x_j, t_1^*)} S(0, t_1^*) \delta(t - t_1^*),$$

if  $(0, t_1) \notin I_+(1)$ .

□

**Claim 4**  $\langle v_j^\varepsilon(x, t), \varphi(x, t) \rangle$  for  $1 \leq j \leq \rho(T)$  converges as  $\varepsilon \rightarrow 0$ .

*Proof.* We prove the claim, using induction on  $j$ . The base case of  $j = 1$  is given by Claims 2 and 3. We make the following assumptions for  $j \leq i - 1$ .

*Assumption 1.*  $\langle v_j^\varepsilon(x, t), \varphi(x, t) \rangle$  converges as  $\varepsilon \rightarrow 0$ .

*Assumption 2.*  $\lim_{\varepsilon \rightarrow 0} v_j^\varepsilon(0, t) = C_j \delta(t - t_j^*)$  in  $\mathcal{D}'(\mathbb{R}_+)$ , where  $C_j$  is a real constant.

Prove the claim for  $j = i$ . Similarly to Claim 2, we distinguish two cases.

*Case 1.*  $(0, t_j) \in I_+(i)$  for some  $j \leq l$ . The claim follows similarly to the proof of Claim 2 for Case 1.

*Case 2.*  $(0, t_j) \notin I_+(i)$ , for  $j \leq l$ . We use (105), (21), and (22), and represent  $v_i^\varepsilon$  in the form

$$v_i^\varepsilon(x, t) = \left[ c_r(\tau) \int_{\omega(\tau; 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon_0))}^{Q(\tau)} b_s^\varepsilon(\xi) v^\varepsilon(0, \theta(\xi, \tau)) S(\xi, \tau) d\xi \right] \Big|_{\tau=\theta(x, t)} S(x, t) \\ + \left[ c_r(\tau) \int_{Q(\tau)}^L (b_s^\varepsilon v_0^\varepsilon)(\xi, \tau) d\xi \right] \Big|_{\tau=\theta(x, t)} S(x, t), \quad (111)$$

where  $Q(t)$  is defined by (37). The convergence of the second summand follows from Claim 1. We now prove the convergence of the first summand. Consider the action

$$\langle c_r(\theta(x, t)) \int_{\omega(\theta(x, t); 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon))}^{Q(\theta(x, t))} b_s^\varepsilon(\xi) v^\varepsilon(0, \theta(\xi, \theta(x, t))) S(\xi, \theta(x, t)) d\xi S(x, t), \varphi(x, t) \rangle \\ = \int_{I_+^{\varepsilon_0}(i)} \int_{\omega(\theta(x, t); 0, t_{i-1}^* + \varepsilon_{i-1}^+(\varepsilon_0))}^{Q(\theta(x, t))} b_s^\varepsilon(\xi) \left[ c_r(\tau) v^\varepsilon(0, \theta(\xi, \tau)) \right. \\ \left. \times S(\xi, \tau) \right] \Big|_{\tau=\theta(x, t)} S(x, t) \varphi(x, t) d\xi d(x, t)$$

$$\begin{aligned}
&= \sum_{(j,q) \in J(i)} \int_{x_j^* - \varepsilon_0}^{x_j^* + \varepsilon_0} \int_{t_q^* - \varepsilon_q^-(\varepsilon_0)}^{t_q^* + \varepsilon_q^+(\varepsilon_0)} \int_0^{P(\xi,t)} \left[ D(\xi, x, t) \varphi(x, \tilde{\omega}(x; 0, \tilde{\omega}(\xi; 0, t))) \right. \\
&\quad \left. - \chi_{[x_j - \varepsilon, x_j + \varepsilon] \times [t_q^* - \varepsilon_q^-(\varepsilon), t_q^* + \varepsilon_q^+(\varepsilon)]} (\xi, t) D(x_j, x, t_q^*) \varphi(x, \tilde{\omega}(x; 0, t_i^*)) \right] b_s^\varepsilon(\xi) v_q^\varepsilon(0, t) dx dt d\xi \\
&\quad + \sum_{(j,q) \in J(i)} \int_{x_j^* - \varepsilon_0}^{x_j^* + \varepsilon_0} b_s^\varepsilon(\xi) d\xi \int_{t_q^* - \varepsilon_q^-(\varepsilon_0)}^{t_q^* + \varepsilon_q^+(\varepsilon_0)} v_q^\varepsilon(0, t) dt \int_0^{P(\xi,t)} D(x_j, x, t_q^*) \varphi(x, \tilde{\omega}(x; 0, t_i^*)) dx,
\end{aligned}$$

where  $J(i)$  is the set of pairs  $(j, q)$  such that  $\omega(x_j; 0, t_q^*) = t_i^*$ ,

$$P(\xi, t) = \begin{cases} L & \text{if } \theta(L, T) \geq \tilde{\omega}(\xi; 0, t), \\ \omega(T; 0, \tilde{\omega}(\xi; 0, t)) & \text{if } \theta(L, T) < \tilde{\omega}(\xi; 0, t), \end{cases} \quad (112)$$

and

$$\begin{aligned}
D(\xi, x, t) &= \left[ c_r(\theta(x, \tau)) S(\xi, \theta(x, \tau)) S(x, \tau) \right. \\
&\quad \left. \times \left[ (\partial_t \theta)(\xi, \theta(x, \tau)) (\partial_t \theta)(x, \tau) \right]^{-1} \right] \Big|_{\tau = \tilde{\omega}(x; 0, \tilde{\omega}(\xi; 0, t))}.
\end{aligned}$$

It is immediate now that

$$\langle v_i^\varepsilon(x, t), \varphi(x, t) \rangle \xrightarrow{\varepsilon \rightarrow 0} \sum_{(j,q) \in J(i)} C_q \int_0^{P(x_j, t_i^*)} D(x_j, x, t_q^*) \varphi(x, \tilde{\omega}(x; 0, t_i^*)) dx.$$

This convergence is true due to the second induction assumption, the condition (10), the continuity of  $c_r, p, \lambda$ , and  $\varphi$ , and the uniform in  $\varepsilon$  boundedness of  $\int_0^T |v^\varepsilon(0, t)| dt$ , that is proved in Lemma 3.

Take a test function  $\psi(t) \in \mathcal{D}(\mathbb{R}_+)$  and consider the action  $\langle v_i^\varepsilon(0, t), \psi(t) \rangle$ . Computation similar to the above shows that

$$\begin{aligned}
v_i^\varepsilon(0, t) &\xrightarrow{\varepsilon \rightarrow 0} c_r(t_i^*) \sum_{(j,r) \in I(i)} \frac{S(x_j, t_i^*)}{(\partial_t \omega)(0; x_j, t_i^*)} S(0, t_i^*) \delta(t - t_i^*) \\
&\quad + c_r(t_i^*) \sum_{(j,q) \in J(i)} C_q \frac{S(x_j, t_i^*)}{(\partial_t \omega)(x_j; 0, t_i^*)} S(0, t_i^*) \delta(t - t_i^*),
\end{aligned}$$

thereby proving the required convergence.  $\square$

Lemma 4 is proved.

## 8 Proof of Lemma 5

We will need two facts from the theory of distributions.

**Fact 1** ([11], 2.2.1) *If  $u \in \mathcal{D}'(X)$  and every point of  $X$  has a neighborhood on which  $u = 0$ , then  $u \equiv 0$ .*

**Fact 2** ([8], 1.5.1) *If  $(f_j)_{1 \leq j < \infty} \in L^1_{loc}(X)$  is a sequence which converges almost everywhere to a function  $f$ , and there is a function  $g \in L^1_{loc}(X)$  such that  $|f_j| \leq g$  for all  $j$ , then  $f \in L^1_{loc}(X)$  and  $f_j \rightarrow f$  in  $\mathcal{D}'(X)$  as  $j \rightarrow \infty$ .*

Fact 2 extends in an obvious way to families of functions  $(f_\varepsilon)_{\varepsilon > 0} \in L^1_{loc}$ .

To prove the lemma, it suffices to show that the conditions assumed in Fact 1 are fulfilled.

Let us fix an arbitrary  $(x, t) \in \Pi \setminus I_+$  and show that there exists a neighborhood  $Y \subset \Pi \setminus I_+$  of  $(x, t)$  such that the restriction of  $v$  to  $Y$  is equal to 0. We choose  $Y$  such that  $\partial Y \cap I_+ = \emptyset$ . This is possible because  $\Pi \setminus I_+$  is open. We now prove that  $v = 0$  on  $Y$ . By the definition of convergence in  $\mathcal{D}'$ , if  $v = \lim_{\varepsilon \rightarrow 0} v^\varepsilon$  in  $\mathcal{D}'(\Pi)$ , then  $v = \lim_{\varepsilon \rightarrow 0} v^\varepsilon$  in  $\mathcal{D}'(Y)$ . On the account of Lemma 4, it suffices to prove the convergence of  $v^\varepsilon$  to 0 in  $\mathcal{D}'(Y)$ .

Let us check the conditions of Fact 2 for  $v^\varepsilon$  on  $Y$ . The function  $v^\varepsilon$  is in  $L^1_{loc}(Y)$  by Proposition 1. By Lemma 1,  $v^\varepsilon$  converges to 0 pointwise on  $Y$  as  $\varepsilon \rightarrow 0$ . By the conditions imposed on  $\lambda$ , each component of  $\Pi \setminus I_+$  is bounded. Since  $Y$  is included in one of the components,  $Y$  is bounded. Clearly, there exists  $T > 0$  such that  $Y \subset \Pi^T$ . Since  $\bigcap_{\varepsilon > 0} I_+^\varepsilon = I_+$ ,  $\partial Y \cap I_+ = \emptyset$ , and  $\text{supp } v^\varepsilon \subset \overline{I_+^\varepsilon}$  (see (31)), there exists  $\tilde{\varepsilon}$  such that  $v^\varepsilon = 0$  on  $Y$  for all  $\varepsilon \geq \tilde{\varepsilon}$ . By (106), that was proved in Section 6, it follows that  $v^\varepsilon = O\left(\frac{1}{\varepsilon}\right)$  on  $\Pi^T$ . This implies the uniform in  $\varepsilon$  estimate

$$|v^\varepsilon| \leq \frac{C}{\varepsilon}$$

on  $Y$ .

Thus all conditions of Fact 2 hold for  $v^\varepsilon$  on  $Y$ . Therefore  $v^\varepsilon \rightarrow 0$  in  $\mathcal{D}'(Y)$  and  $v = 0$  on  $Y$ .

Since  $(x, t) \in \Pi \setminus I_+$  is arbitrary, the lemma is true by Fact 1.

**Remark** *Lemmas 4 and 5 show that  $v^\varepsilon$  are nets converging to measures concentrated on  $I_+$ . From the construction of  $I_+$  (see Definition 2) it follows that in general the density of singularity curves increases as time progresses.*

## Acknowledgments

I am thankful to the members of the DIANA group for their hospitality during my stay at the Vienna university.

## References

- [1] H. Biagioni. Lecture Notes in Mathematics: A Nonlinear Theory of Generalized Functions. Springer-Verlag, Berlin-Heidelberg-New York 1990.
- [2] Bouzinab, A. and Arino, O.: On the existence and uniqueness for an age-dependent population model with nonlinear growth. Facta Univ., Ser. Math. Inf. **8** (1993), 55–68.



- [3] Busenberg, S. and Iannelli, M.: Separable models in age-dependent population dynamics. *J. Math. Biol.* **22** (1985), 145–173.
- [4] Colombeau, J. F.: *Elementary Introduction to New Generalized Functions* (North-Holland Mathematics Studies: Vol. 113). Elsevier Science Publishers 1985.
- [5] Colombeau, J. F. and Oberguggenberger, M.: On a hyperbolic system with a compatible quadratic term: Generalized solutions, delta waves, and multiplication of distributions. *Comm. Part. Diff. Eqs.* **15** (1990), 905–938.
- [6] Demengel, F. and Rauch, J.: Measure valued solutions of asymptotically homogeneous semilinear hyperbolic systems in one space dimension. *Proc. Edinburgh Math. Soc.* **33** (1990), 443–460.
- [7] Demengel, F. and Rauch, J.: Weak convergence of asymptotically homogeneous functions of measures. *Nonlinear Anal.* **15**(1990), 1 - 16.
- [8] Friedlander, G and Joshi, M.: *Introduction to the Theory of Distributions*, Cambridge University Press 1982, second edition 1998.
- [9] Gramchev, T.: Semilinear hyperbolic systems with singular initial data. *Monatsh. Math.* **112** (1991), 99–113.
- [10] Gramchev, T.: Nonlinear maps in spaces of distributions. *Math. Zeitschr.* **209**(1992), 101 - 114.
- [11] Hörmander, L.: *The Analysis of Linear Partial Differential Operators*, Vol. I. Springer 1985, 2nd printing 1994.
- [12] Huyer, W.: Semigroup formulation and approximation of a linear age-dependent population problem with spatial diffusion. *Semigroup Forum* **49** (1994), 99–114.
- [13] Kmit, I. and Hörmann, G.: Semilinear hyperbolic systems with singular non-local boundary conditions: reflection of singularities and delta waves. *J. for Analysis and its Applications* **20** (2001), No. 3, 637–659.
- [14] Oberguggenberger, M.: *Multiplication of distributions and applications to partial differential equations* (Pitman Research Notes in Mathematics: Vol. 259). Harlow: Longman 1992.
- [15] Oberguggenberger, M.: The Carleman system with positive measures as initial data. *Transport Theory Statist. Phys.* **20** (1991), 177–197.
- [16] Oberguggenberger, M.: Weak limits of solutions to semilinear hyperbolic systems. *Math. Ann.* **274** (1986), 599–607.
- [17] Oberguggenberger, M. and Wang, Y. G.: Delta-waves for Semilinear Hyperbolic Cauchy Problems. *Math. Nachr.* **166** (1994), 317–327 .

- [18] Oberguggenberger, M. and Wang, Y. G.: Reflection of delta-waves for nonlinear wave equations in one space variable. *Nonlinear Analysis* **22** (1994), No. 8, 983–992.
- [19] Oberguggenberger, M. and Wang, Y. G.: Semilinear geometric optics for generalized functions. *Z. Anal. Anwend.* **19** (2000), No. 4, 913–926.
- [20] Rauch, J. and Reed, M.: Nonlinear superposition and absorption of delta waves in one space dimension. *Funct. Anal.* **73** (1987), 152–178.
- [21] Sanchez, David. A.: Linear age-dependent population growth with harvesting. *Bull.Math.Biol.* **40** (1978), 377-385.
- [22] Song, J.: Some developments in mathematical demography and their application to the People’s Republic of China. *Theor.Popul.Biol.* **22** (1982), 382–391.